

Bid Antibody

Catalog # ASC10255

Specification

Bid Antibody - Product Information

Application	WB, IHC, IF
Primary Accession	P55957
Other Accession	AAH36364 , 54673639
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	Bid antibody can be used for detection of Bid by Western blot at 0.5 to 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 10 µg/mL.

Bid Antibody - Additional Information

Gene ID **637**
Other Names
 Bid Antibody: FP497, BH3-interacting domain death agonist, p22 BID, BID, BH3 interacting domain death agonist


Target/Specificity
 BID;

Reconstitution & Storage


Bid antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions


Bid Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Western blot analysis of Bid in mouse lung cell lysates with Bid antibody at (A) 0.5, (B) 1, and (C) 2 µg/mL.



Immunohistochemical staining of mouse lung tissue using Bid antibody at 2 µg/mL.



Bid Antibody - Protein Information**Name** BID**Function**

The major proteolytic product p15 BID allows the release of cytochrome c (By similarity). Isoform 1, isoform 2 and isoform 4 induce ICE-like proteases and apoptosis. Isoform 3 does not induce apoptosis. Counters the protective effect of Bcl-2.

Cellular Location

Cytoplasm. Mitochondrion membrane. Mitochondrion outer membrane.
Note=When uncleaved, it is predominantly cytoplasmic. [BH3-interacting domain death agonist p13]: Mitochondrion membrane {ECO:0000250|UniProtKB:P70444}.
Note=Associated with the mitochondrial membrane. {ECO:0000250|UniProtKB:P70444} [Isoform 3]: Cytoplasm

Tissue Location

Isoform 2 and isoform 3 are expressed in spleen, bone marrow, cerebral and cerebellar cortex. Isoform 2 is expressed in spleen, pancreas and placenta (at protein level). Isoform 3 is expressed in lung, pancreas and spleen (at protein level). Isoform 4 is expressed in lung and pancreas (at protein level)

Bid Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Immunofluorescence of Bid in Mouse Lung cells with Bid antibody at 10 µg/mL.

Bid Antibody - Background

Bid Antibody: Apoptosis plays a major role in normal organism development, tissue homeostasis, and removal of damaged cells. Disruption of this process has been implicated in a variety of diseases such as cancer. The Bcl-2 family of proteins is comprised of critical regulators of apoptosis that can be divided into two classes: those that inhibit apoptosis and those that promote cell death. Bid, a pro-apoptotic Bcl-2 family member, is cleaved by caspase-8 in response to apoptotic signals, exposing the Bcl-2 homology 3 (BH3) domain which is normally buried in the full-length protein. The cleaved complex is myristoylated and translocated to the mitochondrial membrane where it may induce mitochondrial Bax and Bak to oligomerize.

Bid Antibody - References

Lockshin RA, Osborne B, and Zakeri Z. Cell death in the third millennium. *Cell Death Differ.* 2000; 7:2-7.
Cory S, Huang DCS, and Adams JM. The Bcl-2 family: roles in cell survival and oncogenesis. *Oncogene* 2003; 22:8590-607.
Heiser D, Labi V, Erlacher M, et al. The Bcl-2 protein family and its role in the development of neoplastic disease. *Exp. Gerontol.* 2004; 39:1125-35.
Wang K, Yin XM, Chao DT, et al. BID: a novel BH3 domain-only death agonist. *Genes Dev.* 1996; 10:2859-69.