

Beclin-1 Antibody

Catalog # ASC10359

Specification

Beclin-1 Antibody - Product Information

Application	WB, IHC
Primary Accession	Q14457
Other Accession	AAH10276 , 16307457
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	Beclin-1 antibody can be used for the detection of Beclin-1 by Western blot at 0.5 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL.

Beclin-1 Antibody - Additional Information

Gene ID **8678**
Other Names
Beclin-1 Antibody: ATG6, VPS30, beclin1, GT197, Beclin-1, Coiled-coil myosin-like BCL2-interacting protein, beclin 1, autophagy related


Target/Specificity
BECN1;

Reconstitution & Storage


Beclin-1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

Beclin-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Western blot analysis of Beclin-1 in 293 cell lysate with Beclin-1 antibody at (A) 0.5, (B) 1 and (C) 2 µg/mL.



Immunohistochemistry of beclin-1 in rat brain tissue with beclin-1 antibody at 2 µg/mL.

Beclin-1 Antibody - Background

Beclin-1 Antibody: Autophagy, the process of bulk degradation of cellular proteins through an autophagosomic-lysosomal pathway is important for normal growth control and may be defective in tumor cells. Beclin-1, a coiled-coil Bcl-2-interacting protein homologous to the yeast autophagy gene *apg6*, is a mammalian autophagy gene that can inhibit tumorigenesis and is expressed at reduced levels in human breast carcinoma, suggesting that defects in autophagy proteins may contribute to the development or progression of tumors. Bcl-2 can bind to Beclin-1 and inhibit Beclin-1-dependent autophagy in yeast and mammalian cells, suggesting that Bcl-2 functions as an

Beclin-1 Antibody - Protein Information**Name** BECN1**Synonyms** GT197**Function**

Plays a central role in autophagy (PubMed:23184933, PubMed:28445460). Acts as core subunit of the PI3K complex that mediates formation of phosphatidylinositol 3-phosphate; different complex forms are believed to play a role in multiple membrane trafficking pathways: PI3KC3-C1 is involved in initiation of autophagosomes and PI3KC3-C2 in maturation of autophagosomes and endocytosis. Involved in regulation of degradative endocytic trafficking and required for the abscission step in cytokinesis, probably in the context of PI3KC3-C2 (PubMed:20643123, PubMed:20208530, PubMed:26783301). Essential for the formation of PI3KC3-C2 but not PI3KC3-C1 PI3K complex forms. Involved in endocytosis (PubMed:25275521). Protects against infection by a neurovirulent strain of Sindbis virus (PubMed:9765397). May play a role in antiviral host defense.

Cellular Location

Cytoplasm
{ECO:0000250|UniProtKB:O88597,
ECO:0000269|PubMed:19713971,
ECO:0000269|PubMed:21364619}. Golgi
apparatus, trans-Golgi network membrane;
Peripheral membrane protein. Endosome
membrane; Peripheral membrane protein.
Endoplasmic reticulum membrane;
Peripheral membrane protein.
Mitochondrion membrane; Peripheral

anti-autophagy protein as well as an anti-apoptotic protein, which helps maintain autophagy at levels that are more compatible with cell survival rather than cell death.

Beclin-1 Antibody - References

Gozuacik D and Kimchi A. Autophagy as a cell death and tumor suppressor mechanism. *Oncogene*. 2004; 23:2891-906.
Kisen GO, Tessitore L, Costelli P, et al. Reduced autophagic activity in primary rat hepatocellular carcinoma and ascites hepatoma cells. *Carcinogenesis* 1993; 14:2501-5.
Liang XH, Kleeman LK, Jiang HH, et al. Protection against fatal Sindbis virus encephalitis by Beclin, a novel Bcl-2-interacting protein. *J. Virol.* 1998; 72:8586-96.
Kametaka S, Okano T, Ohsumi M, et al. Apg14p and Apg6/Vps30p form a protein complex essential for autophagy in the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.* 1998; 273:22284-91.

membrane protein. Endosome
{ECO:0000250|UniProtKB:O88597}
Cytoplasmic vesicle, autophagosome.
Note=Interaction with ATG14 promotes
translocation to autophagosomes.
Expressed in dendrites and cell bodies of
cerebellar Purkinje cells (By similarity)
{ECO:0000250|UniProtKB:O88597,
ECO:0000269|PubMed:19050071}
[Beclin-1-C 37 kDa]: Mitochondrion
{ECO:0000250|UniProtKB:O88597}

Tissue Location

Ubiquitous.

Beclin-1 Antibody - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)