

RIPK1 Antibody

Catalog # ASC10976

Specification

RIPK1 Antibody - Product Information

Application	WB, IHC, IF
Primary Accession	Q13546
Other Accession	NP_003795 , 57242761
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	RIPK1 antibody can be used for detection of RIPK1 by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL.

RIPK1 Antibody - Additional Information

Gene ID **8737**
Target/Specificity
RIPK1;

Reconstitution & Storage


RIPK1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions


RIPK1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RIPK1 Antibody - Protein Information


Name RIPK1 ([HGNC:10019](#))



Western blot analysis of RIPK1 in rat kidney tissue lysate with RIPK1 antibody at 1 µg/mL.



Immunohistochemistry of RIPK1 in mouse kidney tissue with RIPK1 antibody at 2.5 µg/mL.



Immunofluorescence of RIPK1 in Mouse Kidney cells with RIPK1 antibody at 20 µg/mL.

Function

Serine-threonine kinase which is a key regulator of TNF- mediated apoptosis, necroptosis and inflammatory pathways (PubMed:31827280, PubMed:31827281). Exhibits kinase activity-dependent functions that regulate cell death and kinase-independent scaffold functions regulating inflammatory signaling and cell survival (PubMed:11101870, PubMed:19524512, PubMed:19524513, PubMed:29440439, PubMed:30988283). Has kinase-independent scaffold functions: upon binding of TNF to TNFR1, RIPK1 is recruited to the TNF-R1 signaling complex (TNF-RSC also known as complex I) where it acts as a scaffold protein promoting cell survival, in part, by activating the canonical NF-kappa-B pathway (By similarity). Kinase activity is essential to regulate necroptosis and apoptosis, two parallel forms of cell death: upon activation of its protein kinase activity, regulates assembly of two death-inducing complexes, namely complex IIa (RIPK1-FADD-CASP8), which drives apoptosis, and the complex IIb (RIPK1-RIPK3-MLKL), which drives necroptosis (By similarity). RIPK1 is required to limit CASP8- dependent TNFR1-induced apoptosis (By similarity). In normal conditions, RIPK1 acts as an inhibitor of RIPK3-dependent necroptosis, a process mediated by RIPK3 component of complex IIb, which catalyzes phosphorylation of MLKL upon induction by ZBP1 (PubMed:19524512, PubMed:19524513, PubMed:19524513, PubMed:19524513).

RIPK1 Antibody - Background

RIPK1 Antibody: RIPK1 (Receptor Interacting Protein) is a crucial 74 kD adaptor kinase in several of stress-induced signaling pathways and on the crossroad of a cell's decision to live or die. RIPK1 contains an N-terminal region with homology to protein kinases, an intermediate domain capable of association with MAPKKK and a C-terminal region containing a death domain motif present in the Fas and TNFR1 intracellular domains. Full length RIPK1 is important for signalling to NFκ-B, MAPKs and necrosis, whereas caspase-8 generates a C-terminal RIPK1 cleavage fragment, promoting TNF-induced apoptosis. It is required for TNFRSF1A-mediated and TLR3-induced NF-κB activation. RIPK1-deficient mice fail to thrive, displaying extensive apoptosis in both lymphoid and adipose tissues and dying at 1-3 days of age.

RIPK1 Antibody - References

Stanger BZ, Leder P, Lee TH, et al. RIP: a novel protein containing a death domain that interacts with Fas/APO-1 (CD95) in yeast and causes cell death. Cell1995; 81:513-23.
Hsu H, Huang J, Shu HB, et al. TNF-dependent recruitment of the protein kinase RIP to the TNF receptor-1 signaling complex. Immunity1996; 4:387-96.
Meylan E, Burns K, Hofmann K, et al. RIP1 is an essential mediator of Toll-like receptor 3-induced NF-kappa B activation. Nat. Immunol.2004; 5:503-7.
Festjens N, Vanden Bergh T, Cornelis S, et al. RIP1, a kinase on the crossroads of a cell's decision to live or die. Cell Death Differ.2007;14:400-10.

tations/29440439"
target="_blank">29440439,
PubMed:<a href="http://www.uniprot.org/ci
tations/30988283"
target="_blank">30988283). Inhibits
RIPK3- mediated necroptosis via
FADD-mediated recruitment of CASP8,
which cleaves RIPK1 and limits TNF-induced
necroptosis (PubMed:<a href="http://www.
uniprot.org/citations/19524512"
target="_blank">19524512,
PubMed:<a href="http://www.uniprot.org/ci
tations/19524513"
target="_blank">19524513,
PubMed:<a href="http://www.uniprot.org/ci
tations/29440439"
target="_blank">29440439,
PubMed:<a href="http://www.uniprot.org/ci
tations/30988283"
target="_blank">30988283). Required
to inhibit apoptosis and necroptosis during
embryonic development: acts by preventing
the interaction of TRADD with FADD thereby
limiting aberrant activation of CASP8 (By
similarity). In addition to apoptosis and
necroptosis, also involved in inflammatory
response by promoting transcriptional
production of pro-inflammatory cytokines,
such as interleukin-6 (IL6) (PubMed:<a href
="http://www.uniprot.org/citations/3182728
0" target="_blank">31827280,
PubMed:<a href="http://www.uniprot.org/ci
tations/31827281"
target="_blank">31827281).
Phosphorylates RIPK3: RIPK1 and RIPK3
undergo reciprocal auto- and trans-
phosphorylation (PubMed:<a href="http://w
ww.uniprot.org/citations/19524513"
target="_blank">19524513).
Phosphorylates DAB2IP at 'Ser-728' in a
TNF-alpha-dependent manner, and thereby
activates the MAP3K5-JNK apoptotic
cascade (PubMed:<a href="http://www.unip
rot.org/citations/17389591"
target="_blank">17389591,
PubMed:<a href="http://www.uniprot.org/ci
tations/15310755"
target="_blank">15310755). Required
for ZBP1-induced NF-kappa-B activation in
response to DNA damage (By similarity).

Cellular Location

Cytoplasm

{ECO:0000250|UniProtKB:Q60855}. Cell
membrane

{ECO:0000250|UniProtKB:Q9ZUF4}

RIPK1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)