

# KPNA2 Antibody Catalog # ASC11206

# **Specification**

## **KPNA2 Antibody - Product Information**

Application WB, IHC, IF Primary Accession Other Accession NP\_002257, 4504897

Reactivity Human, Mouse,

Host Rabbit
Clonality Polyclonal
Isotype IgG

Application Notes KP

KPNA2 antibody can be used for detection of KPNA2 by

Western blot at 1

- 2 μg/mL.
Antibody can also be used for immu nohistochemistry starting at 5 μg/mL. For immun

ofluorescence start at 20 μg/mL.

# **KPNA2 Antibody - Additional Information**

Gene ID 3838 Target/Specificity KPNA2;


#### **Reconstitution & Storage**

KPNA2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.


### **Precautions**

KPNA2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**KPNA2 Antibody - Protein Information** 



Western blot analysis of KPNA2 in rat heart tissue lysate with KPNA2 antibody at (A) 1 and (B) 2 µg/mL.



Immunohistochemistry of KPNA2 in human heart tissue with KPNA2 antibody at 5  $\mu$ g/mL.



### Name KPNA2 (HGNC:6395)

## Synonyms RCH1, SRP1

#### **Function**

Functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1. Binds specifically and directly to substrates containing either a simple or bipartite NLS motif. Docking of the importin/substrate complex to the nuclear pore complex (NPC) is mediated by KPNB1 through binding to nucleoporin FxFG repeats and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to importin-beta and the three components separate and importin-alpha and -beta are re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran from importin. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus.


Cellular Location Cytoplasm. Nucleus

**Tissue Location**Expressed ubiquitously.

## **KPNA2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



Immunofluorescence of KPNA2 in human heart tissue with KPNA2 antibody at 20 µg/mL.


#### **KPNA2 Antibody - Background**

KPNA2 Antibody: Karvopherin, a cytosolic and heterodimeric protein complex consisting of alpha and beta subunits, is responsible for targeting proteins with nuclear localization signals to the nuclear pore complex by an energy requiring, Ran-dependent mechanism. The alpha subunit and imported substrate enter the nucleus and accumulate in the nucleoplasm, while the beta subunit accumulates at the NPC. KPNA2 is the alpha subunit 2 of karyopherin, which forms a complex with importin subunit beta-1 and functions as a cargo carrier that transports various complexes from cytoplasm into nucleus. It is ubiquitously expressed and contains an IBB/importin beta domain, ten Armadillo repeats that bind "cargo" and three intervening nuclear localization sequences (NLSs). It has recently been reported to play an important role in tumorigenesis and tumor progression.

### **KPNA2 Antibody - References**

Moroianu J. Molecular mechanisms of nuclear protein transport. Crit. Rev. Eukaryot. Gene Expr.1997; 7:61-72.

Gilchrist D and Rexach M. Molecular basis for the rapid dissociation of nuclear localization signals from karyopherin alpha in the nucleoplasm. J. Biol. Chem.2003; 278:





Tel: 858.875.1900 Fax: 858.622.0609

51937-49.

Tseng SF, Chang CY, Wu KJ, et al. Importin KPNA2 is required for proper nuclear localization and multiple functions of NBS1. J. Biol. Chem.2005; 280:39594-600. Sakai M, Sohda M, Miyazaki T, et al. Significance of karyopherin-{alpha} 2 (KPNA2) expression in esophageal squamous cell carcinoma. Anticancer Res.2010; 30:851-6.