

PHOX2B Antibody
 Catalog # **ASC11501**

Specification

PHOX2B Antibody - Product Information

Application	WB
Primary Accession	O99453
Other Accession	NP_003915 , 12707580
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	PHOX2B antibody can be used for detection of PHOX2B by Western blot at 1 - 2 µg/mL.

PHOX2B Antibody - Additional Information

Gene ID **8929**

Target/Specificity

PHOX2B; PHOX2B antibody is predicted to not cross-react with other paired homeobox family members.

Reconstitution & Storage

PHOX2B antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions


PHOX2B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PHOX2B Antibody - Protein Information

Name PHOX2B

Synonyms PMX2B

Function



Western blot analysis of PHOX2B in 293 cell lysate with PHOX2B antibody at (A) 1 and (B) 2 µg/mL.

PHOX2B Antibody - Background

PHOX2B Antibody: PHOX2A and PHOX2B (Paired mesoderm homeobox protein) are closely related, paired-homeodomain transcription factors that function as determinants of the noradrenergic phenotype during embryogenesis. PHOX2 proteins are crucial for the regulation of endogenous hydroxylases in neural crest cells and promote sympathetic neuron generation. Human PHOX2B contains one DNA binding homeobox domain and is required for the differentiation of all central and nonperipheral noradrenergic centers in the brain. In contrast, PHOX2A controls only the differentiation of the main noradrenergic center of the brain. Regulation of PHOX2 may have therapeutic utility in aging or disorders involving degeneration of noradrenergic neurons.

PHOX2B Antibody - References

Stanke M, Junghans D, Geissen M, et al. The Phox2 homeodomain proteins are sufficient to

Involved in the development of several major noradrenergic neuron populations, including the locus coeruleus. Transcription factor which could determine a neurotransmitter phenotype in vertebrates. Enhances second-messenger-mediated activation of the dopamine beta- hydrolase and c-fos promoters, and of several enhancers including cAMP- response element and serum-response element.

Cellular Location

Nucleus

{ECO:0000255|PROSITE-ProRule:PRU00108
}.

Tissue Location

Expressed in neuroblastoma, brain and adrenal gland

promote the development of sympathetic neurons. *Development* 1999; 126:4087-94
Young HM, Ciampoli D, Hsuan J, et al. Expression of Ret-, p75(NTR)-, Phox2a-, Phox2b-, and tyrosine hydroxylase-immunoreactivity by undifferentiated neural crest-derived cells and different classes of enteric neurons in the embryonic mouse gut. *Dev. Dyn.* 1999; 216:137-52.

Pattyn A, Morin X, Cremer H, et al. The homeobox gene Phox2b is essential for the development of autonomic neural crest derivatives. *Nature* 1999; 399:366-70. Coppola E, d'Autréaux F, Rijli FM, et al. Ongoing roles of Phox2 homeodomain transcription factors during neuronal differentiation. *Development* 2010; 137:4211-20

PHOX2B Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)