

CASK Antibody

Catalog # ASC11505

Specification

CASK Antibody - Product Information

Application	WB, IF
Primary Accession	O14936
Other Accession	NP_003679 , 186972120
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	CASK antibody can be used for detection of CASK by Western blot at 1 µg/mL. For immunofluorescence start at 20 µg/mL.

CASK Antibody - Additional Information

Gene ID **8573**
Target/Specificity
 CASK; At least six alternatively spliced transcript variants have been observed.

Reconstitution & Storage

CASK antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.


Precautions

CASK Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


CASK Antibody - Protein Information

Name CASK

Synonyms LIN2



Western blot analysis of CASK in mouse brain tissue lysate with CASK antibody at 1 µg/ml in (A) the absence and (B) the presence of blocking peptide.



Immunofluorescence of CASK in human brain tissue with CASK antibody at 20 µg/mL.

CASK Antibody - Background

CASK Antibody: CASK (Calcium/calmodulin-dependent serine protein kinase), a conserved multi-domain scaffolding protein, belongs to a MAGUK (membrane-associated guanylate kinase

Function

Multidomain scaffolding protein with a role in synaptic transmembrane protein anchoring and ion channel trafficking. Contributes to neural development and regulation of gene expression via interaction with the transcription factor TBR1. Binds to cell-surface proteins, including amyloid precursor protein, neurexins and syndecans. May mediate a link between the extracellular matrix and the actin cytoskeleton via its interaction with syndecan and with the actin/spectrin-binding protein 4.1. Component of the LIN-10-LIN-2-LIN-7 complex, which associates with the motor protein KIF17 to transport vesicles containing N-methyl-D-aspartate (NMDA) receptor subunit NR2B along microtubules (By similarity).

Cellular Location

Nucleus

{ECO:0000250|UniProtKB:Q62915}.

Cytoplasm

{ECO:0000250|UniProtKB:Q62915}. Cell membrane

{ECO:0000250|UniProtKB:Q62915};

Peripheral membrane protein

{ECO:0000250|UniProtKB:Q62915}

Tissue Location

Ubiquitous. Expression is significantly greater in brain relative to kidney, lung, and liver and in fetal brain and kidney relative to lung and liver.

homologs) subfamily and is involved in cell junction organization, tumor suppression and signaling. It is characterized by a novel domain structure that consists of a calcium/calmodulin-dependent protein kinase domain followed by PDZ, SH3 and guanylate kinase-like (GUK) domains. CASK is ubiquitously expressed and significantly greater in brain where it is thought to be involved in signaling at neuronal synapses. CASK interacts with CASKINs and defects in CASK are the cause of mental retardation X-linked CASK-related.

CASK Antibody - References

Dimitratos SD, Woods DF, and Bryant PJ. Camguk, Lin-2, and CASK: novel membrane associated guanylate kinase homologs that also contain CaM kinase domains. *Mech. Dev.* 1997; 63:127-30.

Hsueh YP and Sheng M. Regulated expression and subcellular localization of syndecan heparan sulfate proteoglycans and the syndecan-binding protein CASK/LIN-2 during rat brain development. *J. Neurosci.* 1999;19:7415-25.

Tabuchi K, Biederer T, Butz S, et al. CASK participates in alternative tripartite complexes in which Mint 1 competes for binding with CASKIN2, a novel CASK-binding protein. *J. Neurosci.* 2002; 22:4264-73.

Kuo TY, Hong CJ, Chien HL, et al. X-linked mental retardation gene CASK interacts with Bcl11A/CTIP1 and regulates axon branching and outgrowth. *J. Neurosci. Res.* 2010; 88: 2364-73.

CASK Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)