

N-RAS Antibody Catalog # ASC11665

Specification

N-RAS Antibody - Product Information

Application WB **Primary Accession** P01111 Other Accession NP 002515, 4505451

Reactivity Human, Mouse,

Rat

Host Rabbit Clonality **Polyclonal**

Isotype laG

Calculated MW Predicted: 21

Observed: 20

KDa

Application Notes N-RAS antibody

> can be used for detection of N-RAS by Western

blot at 0.5 - 1

μg/mL.

N-RAS Antibody - Additional Information

Gene ID 4893

Target/Specificity

NRAS; N-RAS antibody is predicted to not cross-react with H-RAS

Reconstitution & Storage

N-RAS antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions


N-RAS Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

N-RAS Antibody - Protein Information

Name NRAS

Synonyms HRAS1

Function



Western blot analysis of N-RAS in 293 cell lysate with N-RAS antibody at (A) 1 and (B) 2 μg/mL.


N-RAS Antibody - Background

N-RAS Antibody: Activating mutations and overexpression of classical Ras subfamily members (N-RAS, H-RAS, and K-RAS) have been widely investigated as key events in the development of human cancers. The N-RAS protein shuttles between the Golgi apparatus and the plasma membrane. This shuttling is regulated through palmitoylation and depalmitoylation by the ZDHHC9-GOLGA7 complex. N-RAS, which has intrinsic GTPase activity, is activated by a quanine nucleotide-exchange factor and inactivated by a GTPase activating protein. Mutations in this gene have been associated with somatic rectal cancer, follicular thyroid cancer, autoimmune lymphoproliferative syndrome, Noonan syndrome, and juvenile myelomonocytic leukemia.

N-RAS Antibody - References

Ehrhardt A, Ehrhardt GR, Guo X, et al. Ras and relatives--job sharing and networking keep an







Ras proteins bind GDP/GTP and possess intrinsic GTPase activity.

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus membrane; Lipid-anchor Note=Shuttles between the plasma membrane and the Golgi apparatus

N-RAS Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cvtometv
- Cell Culture

old family together. Exp. Hematol. 2002; 30:1089-106.

Swarthout JT, Lobo S, Farh L, et al. DHHC9 and GCP9 constitute a human protein fatty acyltransferase with specificity for H- and N-Ras. J. Biol. Chem. 2005; 280:31141-8. Malaney S and Daly RJ. The ras signaling pathway in mammary tumorigenesis and metastasis. J. Mammary Gland Biol. Neoplasia 2001; 6:101-13.

Rodenhuis S. ras and human tumors. Semin Cancer Biol. 1992; 3:241-7