

TMEM107 Antibody

Catalog # ASC11701

Specification

TMEM107 Antibody - Product Information

Application **WB, IHC, IF**
Primary Accession [O6UX40](#)
Other Accession [NP_115730](#),
[34101276](#)
Reactivity **Human**
Host **Rabbit**
Clonality **Polyclonal**
Isotype **IgG**
Calculated MW **Predicted: 16 kDa**

Observed: 19kDa
KDa
Application Notes **TMEM107**
antibody can be
used for
detection of
TMEM107 by
Western blot at 1
- 2 µg/ml.

TMEM107 Antibody - Additional Information

Gene ID **84314**
Target/Specificity
TMEM107; TMEM107 antibody is human specific. At least two isoforms of TMEM107 are known to exist.

Reconstitution & Storage

TMEM107 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions


TMEM107 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TMEM107 Antibody - Protein Information


Name TMEM107 ([HGNC:28128](#))

Function

Plays a role in cilia formation and



Western blot analysis of TMEM107 in THP-1 cell lysate with TMEM107 antibody at 1 µg/ml in (A) the absence and (B) the presence of blocking peptide.




Immunohistochemistry of TMEM107 in human lymph node tissue with TMEM107 antibody at 5 µg/mL.

embryonic patterning. Requires for normal Sonic hedgehog (Shh) signaling in the neural tube and acts in combination with GLI2 and GLI3 to pattern ventral and intermediate neuronal cell types (By similarity). During ciliogenesis regulates the ciliary transition zone localization of some MKS complex proteins (PubMed:26518474).

Cellular Location

Membrane; Multi-pass membrane protein. Cell projection, cilium. Note=Localizes at the transition zone, a region between the basal body and the ciliary axoneme



Immunofluorescence of TMEM107 in human lymph node tissue with TMEM107 antibody at 20 µg/mL.

TMEM107 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TMEM107 Antibody - Background

Transmembrane protein 107 (TMEM 107) is critical for cilia formation and embryonic patterning (1). It is a 139 amino acid transmembrane protein encoded by a gene that maps to human chromosome 17 which is associated with two key tumor suppressor genes p53 and BRCA1 (1,2). TMEM107 is required for normal Sonic hedgehog (Shh) signaling in the neural tube and acts in combination with Gli2 and Gli3 to pattern ventral and intermediate neuronal cell types (1,3). TMEM107 is also a strong candidate gene for central areolar choroidal dystrophy (CACD) (1-3).

TMEM107 Antibody - References

Christopher KJ, Wang B, Kong Y, et al. Forward genetics uncovers transmembrane protein 107 as a novel factor required for ciliogenesis and sonic hedgehog signaling. *Dev. Biol.* 2012; 368:382-92.

Clark HF, Gurney AL, Abaya E, et al. The secreted protein discovery initiative (SPDI), a large-scale effort to identify novel human secreted and transmembrane proteins: a bioinformatics assessment. *Genome Res.* 2003; 13:2265-70.

Ho KS and Scott MP. Sonic hedgehog in the nervous system: functions, modifications and mechanisms. *Curr. Opin. Neurobiol.* 2002; 12:57-63.