

OLIG1 Antibody

Catalog # ASC11782

Specification

OLIG1 Antibody - Product Information

Application **WB, IHC, IF**
Primary Accession [Q8TAK6](#)
Other Accession [NP_620450](#),
[237757328](#)
Reactivity **Human**
Host **Rabbit**
Clonality **Polyclonal**
Isotype **IgG**
Calculated MW **Predicted: 30 kDa**

Observed: 29 kDa
KDa
Application Notes **OLIG1 antibody can be used for detection of OLIG1 by Western blot at 1 - 2 µg/ml. Antibody can also be used for Immunohistochemistry at 5 µg/mL. For Immunofluorescence start at 20 µg/mL.**

OLIG1 Antibody - Additional Information

Gene ID **116448**
Target/Specificity
OLIG1; OLIG1 antibody is human specific. It is predicted to not cross-react with other members of the OLIG family of proteins.


Reconstitution & Storage

OLIG1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.


Precautions

OLIG1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

OLIG1 Antibody - Protein Information



Western blot analysis of OLIG1 in A549 cell lysate with OLIG1 antibody at 1 µg/ml.



Immunohistochemistry of OLIG1 in human brain tissue with OLIG1 antibody at 5 µg/mL.

Name OLIG1**Synonyms** BHLHB6, BHLHE21**Function**

Promotes formation and maturation of oligodendrocytes, especially within the brain. Cooperates with OLIG2 to establish the pMN domain of the embryonic neural tube (By similarity).

Cellular Location

Nucleus

{ECO:0000255|PROSITE-ProRule:PRU00981}.


Tissue Location

Expressed in the brain, in oligodendrocytes. Strongly expressed in oligodendrogliomas, while expression is weak to moderate in astrocytomas. Expression in glioblastomas is highly variable.

OLIG1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Immunofluorescence of OLIG1 in human brain tissue with OLIG1 antibody at 20 µg/mL.

OLIG1 Antibody - Background

The oligodendrocyte transcription factors 1 and 2 (OLIG1 and OLIG2, respectively) make up part of basic helix-loop-helix (bHLH) family of transcription factors that are specifically expressed in zones of the neuroepithelium from which oligodendrocyte precursors emerge (1). Both OLIG1 and OLIG2 genes are downstream targets of Sonic hedgehog and are expressed exclusively in the central nervous system (2). OLIG1 is first expressed in the dorsal portion of the p3 progenitor domain of the ventral neural tube while OLIG2 is first observed in the ventral most p3 domain (2). OLIG1 has been shown to be a SMAD cofactor involved in cell motility induced by transforming growth factor-beta (TGF-beta) (3).

OLIG1 Antibody - References

Zhou Q, Wang S, and Anderson DJ. Identification of a novel family of oligodendrocyte lineage-specific basic helix-loop-helix transcription factors. *Neuron* 2000; 25:331-43.

Lu QR, Yuk D, Alberta JA, et al. Sonic Hedgehog-regulated oligodendrocyte lineage genes encoding bHLH proteins in the mammalian central nervous system. *Neuron* 2000; 25:317-29.

Motizuki M, Isogaya K, Miyake K, et al. Oligodendrocyte transcription factor 1 (Olig1) is a Smad cofactor involved in cell motility induced by transforming growth factor-b. *J. Biol. Chem.* 2013; 288:18911-22.

