

## GAPDH Antibody [12D3D5] (biotin)

Catalog # ASC12042

### Specification

#### GAPDH Antibody [12D3D5] (biotin) - Product Information

Application	WB
Primary Accession	<a href="#">P04406</a>
Other Accession	<a href="#">7669492</a> , <a href="#">NP_002037</a> , <a href="#">2597</a>
Reactivity	Human, Mouse, Rat, Rabbit
Host	Mouse
Clonality	Monoclonal
Isotype	IgG
Calculated MW	36053
Application Notes	Biotin-GAPDH antibody can be used for detection of GAPDH by Western blot at 1 - 2 µg/ml.

#### GAPDH Antibody [12D3D5] (biotin) - Additional Information

Gene ID **2597**

#### Other Names

Glyceraldehyde-3-phosphate dehydrogenase, G3PDH, GAPD

#### Precautions

GAPDH Antibody [12D3D5] (biotin) is for research use only and not for use in diagnostic or therapeutic procedures.


#### GAPDH Antibody [12D3D5] (biotin) - Protein Information

#### Name GAPDH

{ECO:0000303|PubMed:2987855, ECO:0000312|HGNC:HGNC:4141}

#### Function

Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively (PubMed:<a href="http://www.uniprot.org/citations/3170



Western blot analysis of GAPDH in multiple cell and tissue lysates with Biotin-GAPDH antibody at 1 µg/ml. Lanes 1-20: 293, A431, A549, Daudi, HeLa, HepG2, Jurkat, K562, MOLT, 3T3, Raji, Ramos, U937, human brain, mouse brain, rat brain, rabbit brain, mouse lung, mouse liver, and rat liver, respectively.

#### GAPDH Antibody [12D3D5] (biotin) - Background

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD), an important energy-yielding step in carbohydrate metabolism. It also is involved in a number of cellular processes such as membrane fusion, phosphotransferase activity, DNA replication and repair, and nuclear RNA export (1). GAPDH also plays a role in different pathologies such as cancer progression, apoptosis, and neuronal diseases such as Alzheimer's and Huntington's disease (2). GAPDH is constitutively expressed at high levels in almost all tissues and cell lines making it ideal for use as a loading control marker in immunoblots.

#### GAPDH Antibody [12D3D5] (biotin) - References

Sirover MA. New nuclear functions of the glycolytic protein, glyceraldehyde-3-phosphate dehydrogenase, in mammalian cells. J. Cell. Biochem. 2005;

585" target="\_blank">3170585</a>, PubMed:<a href="http://www.uniprot.org/citations/11724794" target="\_blank">11724794</a>). Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D- glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate (PubMed:<a href="http://www.uniprot.org/citations/3170585" target="\_blank">3170585</a>, PubMed:<a href="http://www.uniprot.org/citations/11724794" target="\_blank">11724794</a>). Modulates the organization and assembly of the cytoskeleton (By similarity). Facilitates the CHP1- dependent microtubule and membrane associations through its ability to stimulate the binding of CHP1 to microtubules (By similarity). Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes (PubMed:<a href="http://www.uniprot.org/citations/23071094" target="\_blank">23071094</a>). Upon interferon-gamma treatment assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation (PubMed:<a href="http://www.uniprot.org/citations/23071094" target="\_blank">23071094</a>). Also plays a role in innate immunity by promoting TNF-induced NF-kappa-B activation and type I interferon production, via interaction with TRAF2 and TRAF3, respectively (PubMed:<a href="http://www.uniprot.org/citations/23332158" target="\_blank">23332158</a>, PubMed:<a href="http://www.uniprot.org/citations/27387501" target="\_blank">27387501</a>). Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis (By similarity). Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC (By similarity).

#### Cellular Location

95:45-52.;Glyceraldehyde-3-phosphate dehydrogenase, apoptosis, and neurodegenerative diseases. Annu. Rev. Pharmacol. Toxicol. 2005; 45:269-90.;;

Cytoplasm, cytosol. Nucleus  
{ECO:0000250|UniProtKB:P04797}.  
Cytoplasm, perinuclear region. Membrane  
Cytoplasm, cytoskeleton  
{ECO:0000250|UniProtKB:P04797}  
Note=Translocates to the nucleus following  
S-nitrosylation and interaction with SIAH1,  
which contains a nuclear localization signal  
(By similarity). Postnuclear and Perinuclear  
regions (PubMed:12829261)  
{ECO:0000250|UniProtKB:P04797,  
ECO:0000269|PubMed:12829261}

### **GAPDH Antibody [12D3D5] (biotin) - Protocols**

Provided below are standard protocols that you  
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)