

N-terminal Arginylation Antibody

N-terminal Arginylation Antibody, Clone 4A9

Catalog # ASM10171

Specification

N-terminal Arginylation Antibody - Product Information

Application **WB**
Host **Mouse**
Isotype **IgG1**
Clonality **Monoclonal**

Description

Mouse Anti-N-terminal Arginylation
Monoclonal IgG1

Target/Specificity

Specific for N-terminal arginine next to glutamic acid. Does not detect N-terminal arginine next to aspartic acid or internal arginine residues.

Other Names

N-terminal Arginine Antibody, N-terminal Arginylation Antibody, N-terminal Arginylated Antibody, N terminal Arginine Antibody, N terminal Arginylation Antibody, N terminal Arginylated Antibody, Amino-terminal Arginine Antibody, Amino-terminal Arginylation Antibody, Amino-terminal Arginylated Antibody, Amino terminal Arginine Antibody, Amino terminal Arginylation Antibody, Amino terminal Arginylated Antibody

Immunogen

Synthetic N-terminal arginylated peptide conjugated to KLH

Purification

Protein G Purified

Storage **-20°C**


Storage Buffer

PBS pH 7.4, 50% glycerol, 0.9% Sodium Azide

Shipping **Blue Ice or 4°C**
Temperature

Certificate of Analysis

A 1:1000 dilution of SMC-263 was sufficient for detection of N-terminal Arginylation in 0.5 ug of N-terminal Arginine peptide



Western Blot analysis of N-terminal Arginine-BSA showing detection of 67 kDa N-terminal Arginylation protein using Mouse Anti-N-terminal Arginylation Monoclonal Antibody, Clone 4A9 (ASM10171). Lane 1: Molecular Weight Ladder (MW). Lane 2: BSA. Lane 3: RDHKH-BSA. Lane 4: REHKH-BSA. Lane 5: HKH-BSA. Lane 6: HKERD-BSA. Lane 7: HKRRE-BSA. Load: 0.5 µg. Block: 5% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-N-terminal Arginylation Monoclonal Antibody (ASM10171) at 1:1000 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution (Super Signal West Pico) for 5 min in RT. Predicted/Observed Size: 67 kDa. Other Band(s): 250kDa, 150kDa, 75kDa REHKH-BSA.

N-terminal Arginylation Antibody - Background

Protein arginylation is the post-translational addition of arginine to proteins by arginyltransferase ATE1. Arginylation of proteins has been found to play an essential role in physiological pathways during

conjugated to BSA by ECL immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.

Cellular Localization
Endoplasmic Reticulum

**N-terminal Arginylation Antibody -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

embryogenesis and adulthood (1). Arginylation has also been shown to regulate cell stress responses, including ER stress, cytosolic misfolded proteins, and heat stress (2).

**N-terminal Arginylation Antibody -
References**

1. Saha S. and Kashina A. (2011) Dev Biol. 385(1): 1-8.
2. Deka K., et al. (2016) Cell Death Discov. 2: 16074.