

## UVRAG Antibody

Catalog # ASM10494

### Specification

#### UVRAG Antibody - Product Information

Application **ICC/IF, WB**  
Primary Accession [Q9P2Y5](#)  
Other Accession [NP\\_003360.2](#)  
Host **Rabbit**  
Reactivity **Human, Mouse, Rat**  
Clonality **Polyclonal**  
Format **FITC**

#### Description

Rabbit Anti-Human UVRAG Polyclonal

#### Target/Specificity

Predicted molecular weight at ~78.1kDa.  
Observed molecular weights in the 75-90kDa range.

#### Other Names

DHTX Antibody, p63 Antibody, UV radiation resistance associated Antibody, UVRAG\_HUMAN Antibody

#### Immunogen

Synthetic peptide from the C-terminal of human UVRAG

#### Purification

Peptide Affinity Purified

Storage **-20°C**

#### Storage Buffer

PBS, 50% glycerol, 0.09% sodium azide


Shipping **Blue Ice or 4°C**  
Temperature

#### Certificate of Analysis

A 1:1000 dilution of SPC-605 was sufficient for detection of UVRAG on 293T Rapamycin treated lysates using Goat anti-rabbit IgG:HRP as the secondary antibody.

#### Cellular Localization

Endosome | Lysosome | Endoplasmic reticulum | Nucleus | Chromosome | Centromere




Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-UVRAG Polyclonal Antibody (ASM10494). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Rabbit Anti-UVRAG Polyclonal Antibody (ASM10494) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Rabbit ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Late Endosome, Lysosome, Early Endosome. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) UVRAG Antibody (D) Composite.


## UVRAG Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Western blot analysis of Human 293T showing detection of ~70kDa UVRAG protein using Rabbit Anti-UVRAG Polyclonal Antibody (ASM10494). Lane 1: MW Ladder. Lane 2: Human 293T (20 µg). Load: 20 µg. Block: 5% milk + TBST for 1 hour at RT. Primary Antibody: Rabbit Anti-UVRAG Polyclonal Antibody (ASM10494) at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit: HRP at 1:2000 for 1 hour at RT. Color Development: TMB solution for 12 min at RT. Predicted/Observed Size: ~70kDa.



Western blot analysis of Rat Liver showing detection of ~70kDa UVRAG protein using Rabbit Anti-UVRAG Polyclonal Antibody (ASM10494). Lane 1: MW Ladder. Lane 2: Rat Liver (20 µg). Load: 20 µg. Block: 5% milk + TBST for 1 hour at RT. Primary Antibody: Rabbit Anti-UVRAG Polyclonal Antibody (ASM10494) at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit: HRP at

1:2000 for 1 hour at RT. Color Development:  
TMB solution for 12 min at RT.  
Predicted/Observed Size: ~70kDa.

### **UVRAG Antibody - Background**

UVRAG (UV radiation resistance-associated gene) is associated with the Beclin-1/PI3KC3 complex and promotes PI3KC3 enzymatic activity and autophagy, while suppressing proliferation (1). Beclin-1 binding to UVRAG promotes both autophagosome maturation and endocytic trafficking (2). UVRAG is also a potential tumor suppressor protein with frameshift mutations observed in colon and gastric carcinomas (3-4). It is highly expressed in the brain, lung, kidney and liver.

### **UVRAG Antibody - References**

1. Liang, C. et al. (2008) Nat Cell Biol. 10: 776-87.
2. Ionov, Y. et al. (2004) Oncogene. 23: 639-45.
3. Kim, M.S. et al. (2008) Hum Pathol. 39: 1059-63.