



Fab Anti-Rat IgG (H&L) (Fluorescein Conjugated) Secondary Antibody

Goat Polyclonal, Fluorescein (FITC) Catalog # ASR1725

Specification

Fab Anti-Rat IgG (H&L) (Fluorescein Conjugated) **Secondary Antibody - Product Information**

Description F(ab) Anti-RAT IgG (H&L) (GOAT)

Antibody Fluorescein Conjugated

Host Goat

Conjugate Fluorescein

(FITC)

FP Value 4.0 moles

> Fluorescein (FITC) per mole of IgG

F(ab)

Target Species Rat

Clonality **Polyclonal** Application

,3,4, **Application Note**

FLISA 1:10,000-1:

50,000;IF

Microscopy 1:1,00 0-1:5,000;FlowCyt

ometry

1:500-1:2,500

Physical State Lyophilized Host Isotype IgG F(ab) Target Isotype IgG (H&L) Buffer 0.02 M

Potassium Phosphate, 0.15

M Sodium

Chloride, pH 7.2

Rat IgG whole **Immunogen** molecule

Reconstitution 1.0 mL

Volume

Reconstitution

Restore with Buffer deionized water

(or equivalent)

Stabilizer 10 mg/mL Bovine

Serum Albumin

(BSA) -

Immunoglobulin and Protease

free

0.01% (w/v)Preservative

Sodium Azide

Fab Anti-Rat IgG (H&L) (Fluorescein Conjugated) Secondary Antibody -**Background**

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.





Fab Anti-Rat IgG (H&L) (Fluorescein Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and anti-Goat Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Fab Anti-Rat IgG (H&L) (Fluorescein Conjugated) Secondary Antibody - Protein Information

Fab Anti-Rat IgG (H&L) (Fluorescein Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture