

ACTA2 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a full length recombinant ACTA2.

Catalog # AT1030a

Specification

ACTA2 Antibody (monoclonal) (M02) - Product Information

Application	IF, WB, E
Primary Accession	P62736
Other Accession	BC017554
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b Kappa
Calculated MW	42009

ACTA2 Antibody (monoclonal) (M02) - Additional Information

Gene ID 59

Other Names

Actin, aortic smooth muscle, Alpha-actin-2,
Cell growth-inhibiting gene 46 protein,
ACTA2, ACTSA, ACTVS

Target/Specificity

ACTA2 (AAH17554.1, 1 a.a. ~ 377 a.a)
full-length recombinant protein with GST
tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate
buffered saline, pH 7.2 .


Storage

Store at -20°C or lower. Aliquot to avoid
repeated freezing and thawing.


Precautions

ACTA2 Antibody (monoclonal) (M02) is for
research use only and not for use in
diagnostic or therapeutic procedures.

ACTA2 Antibody (monoclonal) (M02) - Protocols




Immunofluorescence of monoclonal antibody to ACTA2 on HeLa cell. [antibody concentration 1 ~ 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (67.21 KDa) .

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



ACTA2 monoclonal antibody (M01), clone 4A8-2H3. Western Blot analysis of ACTA2 expression in human omentum, serous carcinoma.



Detection limit for recombinant GST tagged ACTA2 is approximately 1ng/ml as a capture antibody.

ACTA2 Antibody (monoclonal) (M02) - Background

The protein encoded by this gene belongs to the actin family of proteins, which are highly conserved proteins that play a role in cell motility, structure and integrity. Alpha, beta and gamma actin isoforms have been identified, with alpha actins being a major constituent of the contractile apparatus, while beta and gamma actins are involved in the regulation of cell motility. This actin is an alpha actin that is found in skeletal muscle. Defects in this gene cause aortic aneurysm familial thoracic type 6. Multiple alternatively spliced variants, encoding the same protein, have been identified.

ACTA2 Antibody (monoclonal) (M02) - References

1.Isolation and Characterization of a Primary Proximal Tubular Epithelial Cell Model from Human Kidney by CD10/CD13 Double Labeling.Van der Hauwaert C, Savary G, Gnemmi V, Glowacki F, Pottier N, Bouillez A, Maboudou P, Zini L, Leroy X, Cauffiez C, Perrais M, Aubert S *PLoS One*. 2013 Jun

14;8(6):e66750. doi:

10.1371/journal.pone.0066750. Print

2013.2.miR-199a-5p Is Upregulated during

Fibrogenic Response to Tissue Injury and

Mediates TGFbeta-Induced Lung Fibroblast

Activation by Targeting Caveolin-1.Lino

Cardenas CL, Henaoui IS, Courcot E, Roderburg

C, Cauffiez C, Aubert S, Copin MC, Wallaert B,

Glowacki F, Dewaeles E, Milosevic J, Maurizio J,

Tedrow J, Marcet B, Lo-Guidice JM, Kaminski N,

Barbry P, Luedde T, Perrais M, Mari B, Pottier

NPLoS Genet. 2013 Feb;9(2):e1003291. doi:

10.1371/journal.pgen.1003291. Epub 2013 Feb

14.