

## AKT1 Antibody (monoclonal) (M08)

Mouse monoclonal antibody raised against a full length recombinant AKT1.

Catalog # AT1098a

### Specification

#### AKT1 Antibody (monoclonal) (M08) - Product Information

|                   |                          |
|-------------------|--------------------------|
| Application       | IF, WB, E                |
| Primary Accession | <a href="#">P31749</a>   |
| Other Accession   | <a href="#">BC000479</a> |
| Reactivity        | Human, Mouse             |
| Host              | mouse                    |
| Clonality         | Monoclonal               |
| Isotype           | IgG2a                    |
| Calculated MW     | 55686                    |

#### AKT1 Antibody (monoclonal) (M08) - Additional Information

Gene ID 207

#### Other Names

RAC-alpha serine/threonine-protein kinase,  
Protein kinase B, PKB, Protein kinase B  
alpha, PKB alpha, Proto-oncogene c-Akt,  
RAC-PK-alpha, AKT1, PKB, RAC

#### Target/Specificity

AKT1 (AAH00479.1, 1 a.a. ~ 480 a.a)  
full-length recombinant protein with GST  
tag. MW of the GST tag alone is 26 KDa.

#### Dilution

WB~~1:500~1000

#### Format

Clear, colorless solution in phosphate  
buffered saline, pH 7.2 .


#### Storage

Store at -20°C or lower. Aliquot to avoid  
repeated freezing and thawing.


#### Precautions

AKT1 Antibody (monoclonal) (M08) is for  
research use only and not for use in  
diagnostic or therapeutic procedures.

#### AKT1 Antibody (monoclonal) (M08) -



Immunofluorescence of monoclonal antibody to AKT1 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (78.54 KDa) .


## Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)




AKT1 monoclonal antibody (M08), clone 1D4.  
Western Blot analysis of AKT1 expression in  
Raw 264.7((Cat # AT1098a )



AKT1 monoclonal antibody (M08), clone 1D4  
Western Blot analysis of AKT1 expression in  
Jurkat ( (Cat # AT1098a )



AKT1 monoclonal antibody (M08), clone 1D4.  
Western Blot analysis of AKT1 expression in  
NIH/3T3((Cat # AT1098a )



Detection limit for recombinant GST tagged AKT1 is approximately 1ng/ml as a capture antibody.

### **AKT1 Antibody (monoclonal) (M08) - Background**

The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Multiple alternatively spliced transcript variants have been found for this gene.

### **AKT1 Antibody (monoclonal) (M08) - References**

HER-2/AKT expression in upper urinary tract urothelial carcinoma: prognostic implications. Izquierdo L, et al. Anticancer Res, 2010 Jun. PMID 20651405. Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. miR-149\* induces apoptosis by inhibiting Akt1 and E2F1 in human cancer cells. Lin RJ, et al. Mol Carcinog, 2010 Aug. PMID 20623644. Genetic variation in a metabolic

signaling pathway and colon and rectal cancer risk: mTOR, PTEN, STK11, RPKAA1, PRKAG2, TSC1, TSC2, PI3K and Akt1. Slattery ML, et al. Carcinogenesis, 2010 Sep. PMID 20622004. The clinicopathological and prognostic relevance of pyruvate kinase M2 and pAkt expression in breast cancer. Benesch C, et al. Anticancer Res, 2010 May. PMID 20592362.