

CACYBP Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant CACYBP. Catalog # AT1370a

Specification

CACYBP Antibody (monoclonal) (M01) - Product Information

Application WB, E **Primary Accession 09HB71** Other Accession BC005975 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IqG1 Kappa Calculated MW 26210

CACYBP Antibody (monoclonal) (M01) - Additional Information

Gene ID 27101

Other Names

Calcyclin-binding protein, CacyBP, hCacyBP, S100A6-binding protein, Siah-interacting protein, CACYBP, S100A6BP, SIP

Target/Specificity

CACYBP (AAH05975, 1 a.a. ~ 228 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.


Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


Precautions

CACYBP Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

CACYBP Antibody (monoclonal) (M01) - Protocols



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (50.82 KDa) .



Western Blot analysis of CACYBP expression in transfected 293T cell line by CACYBP monoclonal antibody (M01), clone 2E3.

Lane 1: CACYBP transfected lysate(26.61 KDa).


Lane 2: Non-transfected lysate.






Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



Detection limit for recombinant GST tagged CACYBP is approximately 1ng/ml as a capture antibody.




Proximity Ligation Analysis of protein-protein interactions between SKP1 and CACYBP. HeLa cells were stained with anti-SKP1 rabbit purified polyclonal 1:1200 and anti-CACYBP mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

CACYBP Antibody (monoclonal) (M01) - Background

The protein encoded by this gene is a calcyclin binding protein. It may be involved in calcium-dependent ubiquitination and subsequent proteosomal degradation of target proteins. It probably serves as a molecular bridge in ubiquitin E3 complexes and participates in the ubiquitin-mediated degradation of beta-catenin. Two alternatively spliced transcript variants encoding different isoforms have been found for this gene.

CACYBP Antibody (monoclonal) (M01) - References

Defining the human deubiquitinating enzyme





interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732. Expression and clinical significance of CacyBP/SIP in pancreatic cancer. Chen X, et al. Pancreatology, 2008. PMID 18765951.Expression of calcyclin-binding protein/Siah-1 interacting protein in normal and malignant human tissues: an immunohistochemical survey. Zhai H, et al. J Histochem Cytochem, 2008 Aug. PMID 18443365. Calcyclin-binding protein inhibits proliferation, tumorigenicity, and invasion of gastric cancer. Ning X, et al. Mol Cancer Res, 2007 Dec. PMID 18171983. Overexpressed CacyBP/SIP leads to the suppression of growth in renal cell carcinoma. Sun S, et al. Biochem Biophys Res Commun, 2007 May 18. PMID 17400182.