

#### CALCA Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant CALCA. Catalog # AT1371a

#### **Specification**

### CALCA Antibody (monoclonal) (M01) - Product Information

Application WB, E **Primary Accession** P06881 Other Accession NM 001741 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2a Kappa Calculated MW 13899

CALCA Antibody (monoclonal) (M01) - Additional

#### Gene ID 796

Information

#### **Other Names**

Calcitonin gene-related peptide 1, Alpha-type CGRP, Calcitonin gene-related peptide I, CGRP-I, CALCA, CALC1

#### Target/Specificity

CALCA (NP\_001732, 52 a.a. ~ 141 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

#### Dilution

WB~~1:500~1000

#### **Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2.


#### **Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


#### **Precautions**

CALCA Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

## CALCA Antibody (monoclonal) (M01) - Protocols



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (35.64 KDa) .



Western Blot analysis of CALCA expression in transfected 293T cell line by CALCA monoclonal antibody (M01), clone 4B10.


Lane 1: CALCA transfected lysate(15.5 KDa). Lane 2: Non-transfected lysate.





Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



Detection limit for recombinant GST tagged CALCA is approximately 1ng/ml as a capture antibody.

# CALCA Antibody (monoclonal) (M01) - Background

This gene encodes the peptide hormones calcitonin, calcitonin gene-related peptide and katacalcin by tissue-specific alternative RNA splicing of the gene transcripts and cleavage of inactive precursor proteins. Calcitonin is involved in calcium regulation and acts to regulate phosphorus metabolism. Calcitonin gene-related peptide functions as a vasodilator while katacalcin is a calcium-lowering peptide. Multiple transcript variants encoding different isoforms have been found for this gene.

### CALCA Antibody (monoclonal) (M01) - References

High-density polymorphisms analysis of 23 candidate genes for association with bone mineral density. Giroux S, et al. Bone, 2010 Jul 30. PMID 20654748. Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Accelerated senescence of endothelial progenitor cells in hypertension is related to the reduction of calcitonin gene-related peptide. Zhou Z, et al. J Hypertens, 2010 May. PMID 20375903.Procalcitonin: a novel cardiac marker with prognostic value in acute coronary syndrome. Atao?lu HE, et al. | Int Med Res, 2010 Jan-Feb. PMID 20233513. Procalcitonin as a prognostic marker in patients with acute myocardial infarction. Kelly D, et al. Biomarkers, 2010 Jun. PMID 20214413.