

CASP10 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant CASP10.

Catalog # AT1402a

Specification

CASP10 Antibody (monoclonal) (M02) - Product Information

Application	IF, WB, E
Primary Accession	Q92851
Other Accession	NM_032974
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	58951

CASP10 Antibody (monoclonal) (M02) - Additional Information

Gene ID 843

Other Names

Caspase-10, CASP-10, Apoptotic protease Mch-4, FAS-associated death domain protein interleukin-1B-converting enzyme 2, FLICE2, ICE-like apoptotic protease 4, Caspase-10 subunit p23/17, Caspase-10 subunit p12, CASP10, MCH4

Target/Specificity

CASP10 (NP_116756, 1 a.a. ~ 110 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format


Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage


Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

CASP10 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.



Immunofluorescence of monoclonal antibody to CASP10 on HeLa cell . [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (37.84 KDa) .


CASP10 Antibody (monoclonal) (M02) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



CASP10 monoclonal antibody (M02), clone 2E7. Western Blot analysis of CASP10 expression in U-2 OS (Cat # AT1402a)



Detection limit for recombinant GST tagged CASP10 is approximately 0.03ng/ml as a capture antibody.

CASP10 Antibody (monoclonal) (M02) - Background

This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 3 and 7, and the protein itself is processed by caspase 8. Mutations in this gene are associated with apoptosis defects seen in type II autoimmune lymphoproliferative syndrome. Three alternatively spliced transcript variants encoding different isoforms have been described for this gene.

CASP10 Antibody (monoclonal) (M02) -

References

Polymorphisms in the caspase genes and the risk of lung cancer. Lee SY, et al. J Thorac Oncol, 2010 Aug. PMID 20661084. A large-scale candidate gene approach identifies SNPs in SOD2 and IL13 as predictive markers of response to preoperative chemoradiation in rectal cancer. Ho-Pun-Cheung A, et al. Pharmacogenomics J, 2010 Jul 20. PMID 20644561. A Large-scale genetic association study of esophageal adenocarcinoma risk. Liu CY, et al. Carcinogenesis, 2010 Jul. PMID 20453000. Mutational analysis of caspase genes in prostate carcinomas. Kim MS, et al. APMIS, 2010 Apr. PMID 20402676. Mutational analysis of CASP10 gene in colon, breast, lung and hepatocellular carcinomas. Oh JE, et al. Pathology, 2010 Jan. PMID 20025484.