

CTH Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a full length recombinant CTH.

Catalog # AT1668a

Specification

CTH Antibody (monoclonal) (M03) - Product Information

Application	WB, IHC, E
Primary Accession	P32929
Other Accession	BC015807
Reactivity	Human, Mouse
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	44508

CTH Antibody (monoclonal) (M03) - Additional Information

Gene ID 1491

Other Names

Cystathionine gamma-lyase,
Cysteine-protein sulphydrase,
Gamma-cystathionase, CTH

Target/Specificity

CTH (AAH15807, 1 a.a. ~ 405 a.a)
full-length recombinant protein with GST
tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate
buffered saline, pH 7.2 .


Storage

Store at -20°C or lower. Aliquot to avoid
repeated freezing and thawing.


Precautions

CTH Antibody (monoclonal) (M03) is for
research use only and not for use in
diagnostic or therapeutic procedures.

CTH Antibody (monoclonal) (M03) - Protocols



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (70.07 KDa) .




CTH monoclonal antibody (M03), clone S51. Western Blot analysis of CTH expression in mouse kidney.

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)




CTH monoclonal antibody (M03), clone S51
Western Blot analysis of CTH expression in K-562 ((Cat # AT1668a)




Western Blot analysis of CTH expression in transfected 293T cell line by CTH monoclonal antibody (M03), clone S51.

Lane 1: CTH transfected lysate(45 KDa).
Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to CTH on formalin-fixed paraffin-embedded human pancreas. [antibody concentration 3

ug/ml]



Detection limit for recombinant GST tagged CTH is approximately 1ng/ml as a capture antibody.

CTH Antibody (monoclonal) (M03) - Background

This gene encodes a cytoplasmic enzyme in the trans-sulfuration pathway that converts cystathione derived from methionine into cysteine. Glutathione synthesis in the liver is dependent upon the availability of cysteine. Mutations in this gene cause cystathioninuria. Alternative splicing of this gene results in three transcript variants encoding different isoforms.

CTH Antibody (monoclonal) (M03) - References

- 1.cGMP-Dependent Protein Kinase Contributes to Hydrogen Sulfide-Stimulated Vasorelaxation.Bucci M, Papapetropoulos A, Vellecco V, Zhou Z, Zaid A, Giannogonas P, Cantalupo A, Dhayade S, Karalis KP, Wang R, Feil R, Cirino G.PLoS One. 2012;7(12):e53319. doi: 10.1371/journal.pone.0053319. Epub 2012 Dec 28.2.Hydrogen sulfide impairs glucose utilization and increases gluconeogenesis in hepatocytes.Zhang L, Yang G, Untereiner A, Ju Y, Wu L, Wang R.Endocrinology. 2013 Jan;154(1):114-26. doi: 10.1210/en.2012-1658. Epub 2012 Nov 26.3.An investigation of the mechanisms of hydrogen sulfide-induced vasorelaxation in rat middle cerebral arteries.Streeter E, Hart J, Badoer E.Naunyn Schmiedebergs Arch Pharmacol. 2012 Jul 17.4.Characterization of Hydrogen Sulfide and Its Synthases, Cystathionine γ -Synthase and Cystathionine γ -Lyase, in Human Prostatic Tissue and Cells.Guo H, Gai JW, Wang Y, Jin HF, Du JB, Jin J.Urology. 2012 Feb;79(2):483.e1-5.5.Glucocorticoids suppress

cystathionine gamma-lyase expression and H₂S production in lipopolysaccharide-treated macrophages. Zhu XY, Liu SJ, Liu YJ, Wang S, Ni X. Cell Mol Life Sci. 2010 Jan 10. [Epub ahead of print]6. H₂S contributes to the hepatic arterial buffer response and mediates vasorelaxation of the hepatic artery via activation of KATP channels. Siebert N, Cantre D, Eipel C, Vollmar B. Am J Physiol Gastrointest Liver Physiol. 2008 Dec;295(6):G1266-73. Epub 2008 Oct 30.