

EGFL7 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a full length recombinant EGFL7.

Catalog # AT1863a

Specification

EGFL7 Antibody (monoclonal) (M02) - Product Information

Application	E
Primary Accession	Q9UHF1
Other Accession	BC012377
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	29618

EGFL7 Antibody (monoclonal) (M02) - Additional Information

Gene ID 51162

Other Names

Epidermal growth factor-like protein 7, EGF-like protein 7, Multiple epidermal growth factor-like domains protein 7, Multiple EGF-like domains protein 7, NOTCH4-like protein, Vascular endothelial statin, VE-statin, Zneu1, EGFL7, MEGF7

Target/Specificity

EGFL7 (AAH12377.1, 24 a.a. ~ 273 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .


Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

EGFL7 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.

EGFL7 Antibody (monoclonal) (M02) - Protocols



Detection limit for recombinant GST tagged EGFL7 is approximately 3ng/ml as a capture antibody.

EGFL7 Antibody (monoclonal) (M02) - References

1.Epidermal Growth Factor-Like Domain 7 Suppresses Intercellular Adhesion Molecule 1 Expression in Response to Hypoxia/Reoxygenation Injury in Human Coronary Artery Endothelial Cells.Badiwala MV, Tumati LC, Joseph JM, Sheshgiri R, Ross HJ, Delgado DH, Rao V.Circulation. 2010 Sep 14;122(11 Suppl):S156-61.

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)