

FASLG Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a partial recombinant FASLG.

Catalog # AT2001a

Specification

FASLG Antibody (monoclonal) (M03) - Product Information

Application	E
Primary Accession	P48023
Other Accession	NM_000639
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	31485

FASLG Antibody (monoclonal) (M03) - Additional Information

Gene ID 356

Other Names

Tumor necrosis factor ligand superfamily member 6, Apoptosis antigen ligand, APTL, CD95 ligand, CD95-L, Fas antigen ligand, Fas ligand, FasL, CD178, Tumor necrosis factor ligand superfamily member 6, membrane form, Tumor necrosis factor ligand superfamily member 6, soluble form, Receptor-binding FasL ectodomain, Soluble Fas ligand, sFasL, ADAM10-processed FasL form, APL, FasL intracellular domain, FasL ICD, SPPL2A-processed FasL form, SPA, FASLG, APT1LG1, CD95L, FASL, TNFSF6

Target/Specificity

FASLG (AAH17502.1, 172 a.a. ~ 281 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Format


Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

FASLG Antibody (monoclonal) (M03) is for



Detection limit for recombinant GST tagged FASLG is approximately 3ng/ml as a capture antibody.

FASLG Antibody (monoclonal) (M03) - Background

The protein encoded by this gene is the ligand for FAS. Both are transmembrane proteins. Interaction of FAS with this ligand is critical in triggering apoptosis of some types of cells such as lymphocytes. Defects in this gene may be related to some cases of systemic lupus erythematosus (SLE).

FASLG Antibody (monoclonal) (M03) - References

Fat intake interacts with polymorphisms of Caspase9, FasLigand and PPARgamma apoptotic genes in modulating Crohn's disease activity. Ferreira P, et al. Clin Nutr, 2010 Jul 20. PMID 20650551. IL-13 and TH2 cytokine exposure triggers matrix metalloproteinase 7-mediated Fas ligand cleavage from bronchial epithelial cells. Wadsworth SJ, et al. J Allergy Clin Immunol, 2010 Aug. PMID 20624652. Physiogenomic analysis of statin-treated patients: domain-specific counter effects within the ACACB gene on low-density lipoprotein cholesterol? Ruafo G, et al. Pharmacogenomics, 2010 Jul. PMID 20602615. Functional polymorphisms in cell death pathway genes and risk of renal cell

research use only and not for use in diagnostic or therapeutic procedures.

FASLG Antibody (monoclonal) (M03) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

carcinoma. Zhu J, et al. Mol Carcinog, 2010 Sep. PMID 20572163. FAS and FASLG genetic variants and risk for second primary malignancy in patients with squamous cell carcinoma of the head and neck. Lei D, et al. Cancer Epidemiol Biomarkers Prev, 2010 Jun. PMID 20501759.