

FBXO22 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant FBXO22.

Catalog # AT2017a

Specification

FBXO22 Antibody (monoclonal) (M01) - Product Information

Application	WB, E
Primary Accession	Q8NEZ5
Other Accession	BC039024
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	44508

FBXO22 Antibody (monoclonal) (M01) - Additional Information

Gene ID 26263

Other Names

F-box only protein 22, F-box protein
FBX22p44, FBXO22, FBX22

Target/Specificity

FBXO22 (AAH39024, 1 a.a. ~ 299 a.a)
full-length recombinant protein with GST
tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate
buffered saline, pH 7.2 .


Storage

Store at -20°C or lower. Aliquot to avoid
repeated freezing and thawing.


Precautions

FBXO22 Antibody (monoclonal) (M01) is for
research use only and not for use in
diagnostic or therapeutic procedures.


FBXO22 Antibody (monoclonal) (M01) - Protocols



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (58.63 KDa) .



FBXO22 monoclonal antibody (M01), clone 6G9 Western Blot analysis of FBXO22 expression in Hela S3 NE ((Cat # AT2017a))



Detection limit for recombinant GST tagged

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FBXO22 is approximately 0.3ng/ml as a capture antibody.

FBXO22 Antibody (monoclonal) (M01) - Background

This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class. Two transcript variants encoding different isoforms exist for this gene.

FBXO22 Antibody (monoclonal) (M01) - References

Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732. FIST: a sensory domain for diverse signal transduction pathways in prokaryotes and ubiquitin signaling in eukaryotes. Borziak K, et al. Bioinformatics, 2007 Oct 1. PMID 17855421. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334. Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039. CUL7: A DOC domain-containing cullin selectively binds Skp1.Fbx29 to form an SCF-like complex. Dias DC, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12481031.