

FH Antibody (monoclonal) (M07)

Mouse monoclonal antibody raised against a full length recombinant FH.

Catalog # AT2044a

Specification

FH Antibody (monoclonal) (M07) - Product Information

Application	IF, WB, E
Primary Accession	P07954
Other Accession	BC003108
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	54637

FH Antibody (monoclonal) (M07) - Additional Information

Gene ID 2271

Other Names

Fumarate hydratase, mitochondrial,
Fumarase, FH

Target/Specificity

FH (AAH03108, 33 a.a. ~ 510 a.a)
full-length recombinant protein with GST
tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate
buffered saline, pH 7.2 .


Storage

Store at -20°C or lower. Aliquot to avoid
repeated freezing and thawing.


Precautions

FH Antibody (monoclonal) (M07) is for
research use only and not for use in
diagnostic or therapeutic procedures.

FH Antibody (monoclonal) (M07) - Protocols




Immunofluorescence of monoclonal antibody to FH on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (78.32 KDa) .

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



FH monoclonal antibody (M07), clone 5C12
Western Blot analysis of FH expression in HeLa (Cat # AT2044a)



Detection limit for recombinant GST tagged FH is approximately 0.1ng/ml as a capture antibody.

FH Antibody (monoclonal) (M07) - Background

The protein encoded by this gene is an enzymatic component of the tricarboxylic acid (TCA) cycle, or Krebs cycle, and catalyzes the formation of L-malate from fumarate. It exists in both a cytosolic form and an N-terminal extended form, differing only in the translation start site used. The N-terminal extended form is targeted to the mitochondrion, where the removal of the extension generates the same form as in the cytoplasm. It is similar to some thermostable class II fumarases and functions as a homotetramer. Mutations in this gene can cause fumarase deficiency and lead to progressive encephalopathy.

FH Antibody (monoclonal) (M07) - References

An approach based on a genome-wide

association study reveals candidate loci for narcolepsy. Shimada M, et al. Hum Genet, 2010 Oct. PMID 20677014. Fumarase: a mitochondrial metabolic enzyme and a cytosolic/nuclear component of the DNA damage response. Yogev O, et al. PLoS Biol, 2010 Mar 9. PMID 20231875. UOK 262 cell line, fumarate hydratase deficient (FH-/FH-) hereditary leiomyomatosis renal cell carcinoma: in vitro and in vivo model of an aberrant energy metabolic pathway in human cancer. Yang Y, et al. Cancer Genet Cytogenet, 2010 Jan 1. PMID 19963135. Follow-up examination of linkage and association to chromosome 1q43 in multiple sclerosis. McCauley JL, et al. Genes Immun, 2009 Oct. PMID 19626040. Novel role of fumarate metabolism in dahl-salt sensitive hypertension. Tian Z, et al. Hypertension, 2009 Aug. PMID 19546378.