

## IL10 Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a partial recombinant IL10.

Catalog # AT2500a

### Specification

#### IL10 Antibody (monoclonal) (M03) - Product Information

|                   |                           |
|-------------------|---------------------------|
| Application       | WB                        |
| Primary Accession | <a href="#">P22301</a>    |
| Other Accession   | <a href="#">NM_000572</a> |
| Reactivity        | Human                     |
| Host              | mouse                     |
| Clonality         | Monoclonal                |
| Isotype           | IgG3 Kappa                |
| Calculated MW     | 20517                     |

#### IL10 Antibody (monoclonal) (M03) - Additional Information

**Gene ID** 3586

#### Other Names

Interleukin-10, IL-10, Cytokine synthesis inhibitory factor, CSIF, IL10

#### Target/Specificity

IL10 (AAI04253.1, 74 a.a. ~ 178 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

#### Dilution

WB~~1:500~1000

#### Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .


#### Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


#### Precautions

IL10 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.

#### IL10 Antibody (monoclonal) (M03) - Protocols



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.29 KDa) .



Western Blot analysis of IL10 expression in transfected 293T cell line by IL10 monoclonal antibody (M03), clone 1C10.

Lane 1: IL10 transfected lysate(20.5 KDa).  
Lane 2: Non-transfected lysate.

#### IL10 Antibody (monoclonal) (M03) - Background

The protein encoded by this gene is a cytokine produced primarily by monocytes and to a

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract.

#### **IL10 Antibody (monoclonal) (M03) - References**

Association of Interleukin-10 Polymorphisms with Cytokines in Type 2 Diabetic Nephropathy. Kung WJ, et al. Diabetes Technol Ther, 2010 Sep 1. PMID 20809684. Interleukin-10 microsatellite variants and the risk of acute coronary syndrome among Tunisians. Ben-Hadj-Khalifa-Kechiche S, et al. Int J Immunogenet, 2010 Aug 29. PMID 20807264. Interleukin-10 promoter polymorphism in mild cognitive impairment and in its clinical evolution. Arosio B, et al. Int J Alzheimers Dis, 2010 Jul 20. PMID 20798845. The associations between two polymorphisms in the interleukin-10 gene promoter and breast cancer risk. Yu KD, et al. Breast Cancer Res Treat, 2010 Aug 25. PMID 20737205. Association of genetic polymorphisms in the interleukin-10 promoter with risk of prostate cancer in Chinese. Liu J, et al. BMC Cancer, 2010 Aug 24. PMID 20735825.