

IL23R Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant IL23R. Catalog # AT2517a

Specification

IL23R Antibody (monoclonal) (M01) - Product Information

Application WB, E **Primary Accession** O5VWK5 Other Accession NM 144701 Reactivity Human Host mouse Clonality Monoclonal Isotype IgG2a Kappa

Calculated MW 71722

IL23R Antibody (monoclonal) (M01) - Additional Information

Gene ID 149233

Other Names

Interleukin-23 receptor, IL-23 receptor, IL-23R, IL23R

Target/Specificity

IL23R (NP 653302, 553 a.a. ~ 628 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.


Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


Precautions

IL23R Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


IL23R Antibody (monoclonal) (M01) -**Protocols**



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (34.1 KDa).



IL23R monoclonal antibody (M01), clone 3D7 Western Blot analysis of IL23R expression in A-431 ((Cat # AT2517a)



Detection limit for recombinant GST tagged





Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

IL23R is approximately 0.1ng/ml as a capture antibody.

IL23R Antibody (monoclonal) (M01) - Background

The protein encoded by this gene is a subunit of the receptor for IL23A/IL23. This protein pairs with the receptor molecule IL12RB1/IL12Rbeta1, and both are required for IL23A signaling. This protein associates constitutively with Janus kinase 2 (JAK2), and also binds to transcription activator STAT3 in a ligand-dependent manner.

IL23R Antibody (monoclonal) (M01) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.Toll-like receptor 9, NOD2 and IL23R gene polymorphisms influenced outcome in AML patients transplanted from HLA-identical sibling donors. Elmaagacli AH, et al. Bone Marrow Transplant, 2010 Jul 12. PMID 20622911.Genome-wide association studies identify IL23R-IL12RB2 and IL10 as Beh?et's disease susceptibility loci. Mizuki N, et al. Nat Genet, 2010 Aug. PMID 20622879. Genome-wide association study identifies variants in the MHC class I, IL10, and IL23R-IL12RB2 regions associated with Beh?et's disease. Remmers EF, et al. Nat Genet, 2010 Aug. PMID 20622878.A nonsynonymous polymorphism in IL23R gene is associated with risk of gastric cancer in a Chinese population. Chen J, et al. Mol Carcinog, 2010 Oct. PMID 20607725.