

MAPK13 Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a partial recombinant MAPK13. Catalog # AT2787a

Specification

MAPK13 Antibody (monoclonal) (M03) - Product Information

Application IF, WB, IHC, E **Primary Accession** 015264 Other Accession BC000433 Reactivity Human Host Mouse Clonality **Monoclonal** Isotype IgG1 Kappa Calculated MW 42090

MAPK13 Antibody (monoclonal) (M03) - Additional Information

Gene ID 5603

Other Names

Mitogen-activated protein kinase 13, MAP kinase 13, MAPK 13, Mitogen-activated protein kinase p38 delta, MAP kinase p38 delta, Stress-activated protein kinase 4, MAPK13, PRKM13, SAPK4

Target/Specificity

MAPK13 (AAH00433, 251 a.a. \sim 365 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage


Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

MAPK13 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.



Immunofluorescence of monoclonal antibody to MAPK13 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (38.28 KDa).




MAPK13 Antibody (monoclonal) (M03) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



MAPK13 monoclonal antibody (M03), clone 1E11 Western Blot analysis of MAPK13 expression in A-549 ((Cat # AT2787a)



Immunoperoxidase of monoclonal antibody to MAPK13 on formalin-fixed paraffin-embedded human small Intestine. [antibody concentration 1 ug/ml]



Detection limit for recombinant GST tagged MAPK13 is approximately 0.3ng/ml as a capture antibody.

MAPK13 Antibody (monoclonal) (M03) - Background

The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as





an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is closely related to p38 MAP kinase, both of which can be activated by proinflammatory cytokines and cellular stress. MAP kinase kinases 3, and 6 can phosphorylate and activate this kinase. Transcription factor ATF2, and microtubule dynamics regulator stathmin have been shown to be the substrates of this kinase.

MAPK13 Antibody (monoclonal) (M03) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Human p38 delta MAP kinase mediates UV irradiation induced up-regulation of the gene expression of chemokine BRAK/CXCL14. Ozawa S, et al. Biochem Biophys Res Commun, 2010 Jun 11. PMID 20478268.p38delta mitogen-activated protein kinase regulates skin homeostasis and tumorigenesis. Efimova T. Cell Cycle, 2010 Feb 1. PMID 20090411. Polymorphisms in innate immunity genes and patients response to dendritic cell-based HIV immuno-treatment. Segat L, et al. Vaccine, 2010 Mar 2. PMID 20056178.Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet. 2009 Nov. PMID 19913121.