

PECAM1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant PECAM1. Catalog # AT3267a

Specification

PECAM1 Antibody (monoclonal) (M01) - Product Information

Application IP, WB, IHC **Primary Accession** P16284 Other Accession BC022512 Reactivity Human Host Mouse Clonality **Monoclonal** Isotype IgG1 kappa Calculated MW 82522

PECAM1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 5175

Other Names

Platelet endothelial cell adhesion molecule, PECAM-1, EndoCAM, GPIIA', PECA1, CD31, PECAM1

Target/Specificity

PECAM1 (AAH22512, 29 a.a. ~ 738 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


Precautions

PECAM1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

PECAM1 Antibody (monoclonal) (M01) - Protocols



Immunoprecipitation of PECAM1 transfected lysate using anti-PECAM1 monoclonal antibody and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with PECAM1 MaxPab rabbit polyclonal antibody.




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (103.95 KDa).






Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture




PECAM1 monoclonal antibody (M01), clone 1D2-1A5. Western Blot analysis of PECAM1 expression in human placenta.



Western Blot analysis of PECAM1 expression in transfected 293T cell line by PECAM1 monoclonal antibody (M01), clone 1D2-1A5.

Lane 1: PECAM1 transfected lysate(82.4 KDa).

Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to PECAM1 on formalin-fixed





paraffin-embedded human lung, adenosquamous cell carcinoma tissue. [antibody concentration 5 ug/ml]

PECAM1 Antibody (monoclonal) (M01) - Background

The protein encoded by this gene is found on the surface of platelets, monocytes, neutrophils, and some types of T-cells, and makes up a large portion of endothelial cell intercellular junctions. The encoded protein is a member of the immunoglobulin superfamily and is likely involved in leukocyte migration, angiogenesis, and integrin activation.

PECAM1 Antibody (monoclonal) (M01) - References

A genetic association study of maternal and fetal candidate genes that predispose to preterm prelabor rupture of membranes (PROM). Romero R, et al. Am J Obstet Gynecol, 2010 Jul 29. PMID 20673868. Physiogenomic analysis of statin-treated patients: domain-specific counter effects within the ACACB gene on low-density lipoprotein cholesterol? Rua?o G, et al. Pharmacogenomics, 2010 Jul. PMID 20602615. Phosphorylation of leukocyte PECAM and its association with detergent-resistant membranes regulate transendothelial migration. Florey O, et al. | Immunol, 2010 Aug 1. PMID 20581150.Interleukin-9 polymorphism in infants with respiratory syncytial virus infection: an opposite effect in boys and girls. Schuurhof A, et al. Pediatr Pulmonol, 2010 Jun. PMID 20503287. Common polymorphisms in ITGA2, PON1 and THBS2 are associated with coronary atherosclerosis in a candidate gene association study of the Chinese Han population. Wang Y, et al. | Hum Genet, 2010 Aug. PMID 20485444.