

PIN1 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant PIN1.

Catalog # AT3313a

Specification

PIN1 Antibody (monoclonal) (M02) - Product Information

Application	WB, E
Primary Accession	Q13526
Other Accession	BC002899
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 kappa
Calculated MW	18243

PIN1 Antibody (monoclonal) (M02) - Additional Information

Gene ID 5300

Other Names

Peptidyl-prolyl cis-trans isomerase
NIMA-interacting 1, Peptidyl-prolyl cis-trans
isomerase Pin1, PPlase Pin1, Rotamase
Pin1, PIN1

Target/Specificity

PIN1 (AAH02899, 64 a.a. ~ 163 a.a) partial
recombinant protein with GST tag. MW of
the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate
buffered saline, pH 7.2 .


Storage

Store at -20°C or lower. Aliquot to avoid
repeated freezing and thawing.


Precautions

PIN1 Antibody (monoclonal) (M02) is for
research use only and not for use in
diagnostic or therapeutic procedures.

PIN1 Antibody (monoclonal) (M02) -



Antibody Reactive Against Recombinant
Protein. Western Blot detection against
Immunogen (36.63 KDa) .



Detection limit for recombinant GST tagged
PIN1 is approximately 0.1ng/ml as a capture
antibody.

PIN1 Antibody (monoclonal) (M02) - Background

Peptidyl-prolyl cis/trans isomerases (PPlases; EC 5.2.1.8), such as PIN1, catalyze the cis/trans isomerization of peptidyl-prolyl peptide bonds. PIN1 is the only PPlase that specifically binds to phosphorylated ser/thr-pro motifs to catalytically regulate the post-phosphorylation conformation of its substrates. PIN1-catalyzed conformational regulation has a profound impact on key proteins involved in the regulation of cell growth, genotoxic and other

Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

stress responses, the immune response, germ cell development, neuronal differentiation, and survival (review by Lu and Zhou, 2007 [PubMed 17878917]).

PIN1 Antibody (monoclonal) (M02) - References

A PIN1 polymorphism that prevents its suppression by AP4 associates with delayed onset of Alzheimer's disease. Ma SL, et al. Neurobiol Aging, 2010 Jun 24. PMID 20580132. Uncoating of human immunodeficiency virus type 1 requires prolyl isomerase Pin1. Misumi S, et al. J Biol Chem, 2010 Aug 13. PMID 20529865. The prolyl isomerase Pin1 induces LC-3 expression and mediates tamoxifen resistance in breast cancer. Namgoong GM, et al. J Biol Chem, 2010 Jul 30. PMID 20479004. Elevated PIN1 expression by C/EBPalpha-p30 blocks C/EBPalpha-induced granulocytic differentiation through c-Jun in AML. Pulikkan JA, et al. Leukemia, 2010 May. PMID 20376080. Membrane permeable cyclic peptidyl inhibitors against human Peptidylprolyl Isomerase Pin1. Liu T, et al. J Med Chem, 2010 Mar 25. PMID 20180533.