

RNF111 Antibody (monoclonal) (M05)

Mouse monoclonal antibody raised against a partial recombinant RNF111.

Catalog # AT3657a

Specification

RNF111 Antibody (monoclonal) (M05) - Product Information

Application	IF, WB, E
Primary Accession	Q6ZNA4
Other Accession	NM_017610
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	108862

RNF111 Antibody (monoclonal) (M05) - Additional Information

Gene ID 54778

Other Names

E3 ubiquitin-protein ligase Arkadia, 632-, RING finger protein 111, RNF111

Target/Specificity

RNF111 (NP_060080, 1 a.a. ~ 108 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .


Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


Precautions

RNF111 Antibody (monoclonal) (M05) is for research use only and not for use in diagnostic or therapeutic procedures.

RNF111 Antibody (monoclonal) (M05) - Protocols




Immunofluorescence of monoclonal antibody to RNF111 on HeLa cell . [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.62 KDa) .

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)




Western Blot analysis of RNF111 expression in transfected 293T cell line by RNF111 monoclonal antibody (M05), clone 1C4.

Lane 1: RNF111 transfected lysate (107.8 kDa).

Lane 2: Non-transfected lysate.



Western blot analysis of RNF111 over-expressed 293 cell line, cotransfected with RNF111 Validated Chimera RNAi (Cat # AT3657a)



Detection limit for recombinant GST tagged RNF111 is 0.03 ng/ml as a capture antibody.

RNF111 Antibody (monoclonal) (M05) - Background

The protein encoded by this gene contains a RING finger domain, a motif known to be involved in protein-protein and protein-DNA interactions. The mouse counterpart of this gene (Rnf111/arkadia) has been shown to genetically interact with the transforming growth factor (TGF) beta-like factor Nodal, and act as a modulator of the nodal signaling cascade, which is essential for the induction of mesoderm during embryonic development.

RNF111 Antibody (monoclonal) (M05) - References

1. RNF111/Arkadia is a SUMO-targeted ubiquitin ligase that facilitates the DNA damage response. Poulsen SL, Hansen RK, Wagner SA, van Cuijk L, van Belle GJ, Streicher W, Wikstrom M, Choudhary C, Houtsmuller AB, Marteijn JA, Bekker-Jensen S, Mailand NJ Cell Biol. 2013 Jun 10;201(6):797-807. doi: 10.1083/jcb.201212075. 2. Arkadia, a Novel SUMO-Targeted Ubiquitin Ligase Involved in PML Degradation. Erker Y, Neyret-Kahn H, Seeler JS, Dejean A, Atfi A, Levy LMol Cell Biol. 2013 Jun;33(11):2163-77. doi: 10.1128/MCB.01019-12. Epub 2013 Mar 25. 3. RNF111-Dependent Neddylation Activates DNA Damage-Induced Ubiquitination. Ma T, Chen Y, Zhang F, Yang CY, Wang S, Yu XMol Cell. 2013 Mar 7;49(5):897-907. doi: 10.1016/j.molcel.2013.01.006. Epub 2013 Feb 7. 4. RB1CC1 positively regulates transforming growth factor- β signaling through the modulation of Arkadia E3 ubiquitin ligase activity. Koinuma D, Shinozaki M, Nagano Y, Ikushima H, Horiguchi K, Goto K, Chano T, Saitoh M, Imamura T, Miyazono K, Miyazawa K. J Biol Chem. 2011 Jul 27. [Epub ahead of print]. 5. Efficient TGF- β /SMAD signaling in human melanoma cells associated with high c-SKI/SnoN expression. Javelaud D, van Kempen L, Alexaki VI, Le Scolan E, Luo K, Mauviel A. Mol Cancer. 2011 Jan 6;10(1):2. 6. Context-dependent regulation of the expression of c-Ski protein by Arkadia in human cancer cells. Nagano Y, Koinuma D, Miyazawa K, Miyazono K. J Biochem. 2010

Apr;147(4):545-54. Epub 2009 Dec
2.7.Overexpression of snon/skil, amplified at
the 3q26.2 locus, in ovarian cancers: a role in
ovarian pathogenesis.Nanjundan M, Cheng KW,
Zhang F, Lahad J, Kuo WL, Schmandt R,
Smith-McCune K, Fishman D, Gray JW, Mills
GB.Mol Oncol. 2008 Aug;2(2):164-81. Epub
2008 May 10.8.Arkadia activates
Smad3/Smad4-dependent transcription by
triggering signal-induced SnoN
degradation.Levy L, Howell M, Das D, Harkin S,
Episkopou V, Hill CS.Mol Cell Biol. 2007
Sep;27(17):6068-83. Epub 2007 Jun 25.