

RUNX1 Antibody (monoclonal) (M05)

Mouse monoclonal antibody raised against a partial recombinant RUNX1.

Catalog # AT3738a

Specification

RUNX1 Antibody (monoclonal) (M05) - Product Information

Application	IF, WB, IHC, E
Primary Accession	Q01196
Other Accession	NM_001001890
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b Kappa
Calculated MW	48737

RUNX1 Antibody (monoclonal) (M05) - Additional Information

Gene ID 861

Other Names

Runt-related transcription factor 1, Acute myeloid leukemia 1 protein, Core-binding factor subunit alpha-2, CBF-alpha-2, Oncogene AML-1, Polyomavirus enhancer-binding protein 2 alpha B subunit, PEA2-alpha B, PEBP2-alpha B, SL3-3 enhancer factor 1 alpha B subunit, SL3/AKV core-binding factor alpha B subunit, RUNX1, AML1, CBFA2

Target/Specificity

RUNX1 (NP_001001890.1, 210 a.a. ~ 310 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format


Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage


Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

RUNX1 Antibody (monoclonal) (M05) is for



Immunofluorescence of monoclonal antibody to RUNX1 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.85 KDa) .

research use only and not for use in diagnostic or therapeutic procedures.


RUNX1 Antibody (monoclonal) (M05) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)




RUNX1 monoclonal antibody (M05), clone 4E7 Western Blot analysis of RUNX1 expression in HeLa S3 NE (Cat # AT3738a)



RUNX1 monoclonal antibody (M05), clone 4E7. Western Blot analysis of RUNX1 expression in NIH/3T3 (Cat # AT3738a)



Immunoperoxidase of monoclonal antibody to RUNX1 on formalin-fixed paraffin-embedded human placenta. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged RUNX1 is approximately 1ng/ml as a capture antibody.

RUNX1 Antibody (monoclonal) (M05) - Background

Core binding factor (CBF) is a heterodimeric transcription factor that binds to the core element of many enhancers and promoters. The protein encoded by this gene represents the alpha subunit of CBF and is thought to be involved in the development of normal hematopoiesis. Chromosomal translocations involving this gene are well-documented and have been associated with several types of leukemia. Three transcript variants encoding different isoforms have been found for this gene.

RUNX1 Antibody (monoclonal) (M05) - References

Clinical significance of runt-related transcription factor 1 polymorphism in prostate cancer. Huang SP, et al. BJU Int, 2010 Aug 24. PMID 20735389. Therapy-related, mixed phenotype acute leukemia with t(1;21)(p36;q22) and RUNX1 rearrangement. Yamamoto K, et al. Cancer Genet Cytogenet, 2010 Sep. PMID 20682397. Combined mutations of ASXL1, CBL, FLT3, IDH1, IDH2, JAK2, KRAS, NPM1, NRAS, RUNX1, TET2 and WT1 genes in myelodysplastic syndromes and acute myeloid leukemias. Rocquain J, et al. BMC Cancer, 2010 Aug 2. PMID 20678218. Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Genome-wide analysis of estrogen receptor alpha DNA binding and

tethering mechanisms identifies Runx1 as a novel tethering factor in receptor-mediated transcriptional activation. Stender JD, et al. Mol Cell Biol, 2010 Aug. PMID 20547749.