

## SMAD4 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant SMAD4.

Catalog # AT3942a

### Specification

#### SMAD4 Antibody (monoclonal) (M01) - Product Information

|                   |                          |
|-------------------|--------------------------|
| Application       | IF, WB                   |
| Primary Accession | <a href="#">Q13485</a>   |
| Other Accession   | <a href="#">BC002379</a> |
| Reactivity        | Human                    |
| Host              | mouse                    |
| Clonality         | Monoclonal               |
| Isotype           | IgG2a Kappa              |
| Calculated MW     | 60439                    |

#### SMAD4 Antibody (monoclonal) (M01) - Additional Information

Gene ID 4089

#### Other Names

Mothers against decapentaplegic homolog 4, MAD homolog 4, Mothers against DPP homolog 4, Deletion target in pancreatic carcinoma 4, SMAD family member 4, SMAD 4, Smad4, hSMAD4, SMAD4, DPC4, MADH4

#### Target/Specificity

SMAD4 (AAH02379, 1 a.a. ~ 552 a.a)  
full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

#### Dilution

WB~~1:500~1000

#### Format


Clear, colorless solution in phosphate buffered saline, pH 7.2 .

#### Storage


Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

#### Precautions

SMAD4 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.



Immunofluorescence of monoclonal antibody to SMAD4 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (86.46 KDa) .

## SMAD4 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



## SMAD4 Antibody (monoclonal) (M01) - Background

This gene encodes a member of the Smad family of signal transduction proteins. Smad proteins are phosphorylated and activated by transmembrane serine-threonine receptor kinases in response to TGF-beta signaling. The product of this gene forms homomeric complexes and heteromeric complexes with other activated Smad proteins, which then accumulate in the nucleus and regulate the transcription of target genes. This protein binds to DNA and recognizes an 8-bp palindromic sequence (GTCTAGAC) called the Smad-binding element (SBE). The Smad proteins are subject to complex regulation by post-translational modifications. Mutations or deletions in this gene have been shown to result in pancreatic cancer, juvenile polyposis syndrome, and hereditary hemorrhagic telangiectasia syndrome.

## SMAD4 Antibody (monoclonal) (M01) - References

SMAD4 mediates mesenchymal-epithelial reversion in SW480 colon carcinoma cells. Pohl M, et al. Anticancer Res, 2010 Jul. PMID 20682989. Maternal genes and facial clefts in offspring: a comprehensive search for genetic associations in two population-based cleft studies from Scandinavia. Jugessur A, et al. PLoS One, 2010 Jul 9. PMID 20634891. Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril

and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. SMAD4--molecular gladiator of the TGF-beta signaling is trampled upon by mutational insufficiency in colorectal carcinoma of Kashmiri population: an analysis with relation to KRAS proto-oncogene. Sameer AS, et al. BMC Cancer, 2010 Jun 17. PMID 20565773. Expression of oncogenic K-ras and loss of Smad4 cooperate to induce the expression of EGFR and to promote invasion of immortalized human pancreas ductal cells. Zhao S, et al. Int J Cancer, 2010 Nov 1. PMID 20473902.