

SNAP25 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant SNAP25.

Catalog # AT3972a

Specification

SNAP25 Antibody (monoclonal) (M01) - Product Information

Application	WB, E
Primary Accession	P60880
Other Accession	BC010647
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	23315

SNAP25 Antibody (monoclonal) (M01) - Additional Information

Gene ID 6616

Other Names

Synaptosomal-associated protein 25,
SNAP-25, Super protein, SUP,
Synaptosomal-associated 25 kDa protein,
SNAP25, SNAP

Target/Specificity

SNAP25 (AAH10647, 31 a.a. ~ 130 a.a)
partial recombinant protein with GST tag.
MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .


Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


Precautions

SNAP25 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

SNAP25 Antibody (monoclonal) (M01) -



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .



Western Blot analysis of SNAP25 expression in transfected 293T cell line by SNAP25 monoclonal antibody (M01), clone 4A3.


Lane 1: SNAP25 transfected lysate(23.315 KDa).

Lane 2: Non-transfected lysate.

Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Detection limit for recombinant GST tagged SNAP25 is 0.03 ng/ml as a capture antibody.

SNAP25 Antibody (monoclonal) (M01) - Background

Synaptic vesicle membrane docking and fusion is mediated by SNAREs (soluble N-ethylmaleimide-sensitive factor attachment protein receptors) located on the vesicle membrane (v-SNAREs) and the target membrane (t-SNAREs). The assembled v-SNARE/t-SNARE complex consists of a bundle of four helices, one of which is supplied by v-SNARE and the other three by t-SNARE. For t-SNAREs on the plasma membrane, the protein syntaxin supplies one helix and the protein encoded by this gene contributes the other two. Therefore, this gene product is a presynaptic plasma membrane protein involved in the regulation of neurotransmitter release. Two alternative transcript variants encoding different protein isoforms have been described for this gene.

SNAP25 Antibody (monoclonal) (M01) - References

Association Among SNAP-25 Gene Ddel and MnlI Polymorphisms and Hemodynamic Changes During Methylphenidate Use: A Functional Near-Infrared Spectroscopy Study. Oner O, et al. J Atten Disord, 2010 Aug 2. PMID 20679152. An approach based on a genome-wide association study reveals candidate loci for narcolepsy. Shimada M, et al. Hum Genet, 2010 Oct. PMID 20677014. An association study between SNAP-25 gene and Attention-Deficit Hyperactivity Disorder. Zhang H, et al. Eur J Paediatr Neurol, 2010 Jul 2. PMID 20599404. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID

20379614.Association between a synaptosomal protein (SNAP-25) gene polymorphism and verbal memory and attention in patients with endogenous psychoses and mentally healthy subjects. Golimbet VE, et al. Neurosci Behav Physiol, 2010 May. PMID 20333500.