

SPINK1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant SPINK1.

Catalog # AT4023a

Specification

SPINK1 Antibody (monoclonal) (M01) - Product Information

Application	IP, WB, IHC, E
Primary Accession	P00995
Other Accession	BC025790
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	8507

SPINK1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 6690

Other Names

Pancreatic secretory trypsin inhibitor,
Serine protease inhibitor Kazal-type 1,
Tumor-associated trypsin inhibitor, TATI,
SPINK1, PSTI

Target/Specificity

SPINK1 (AAH25790, 24 a.a. ~ 79 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .


Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


Precautions

SPINK1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

SPINK1 Antibody (monoclonal) (M01) -



Immunoprecipitation of SPINK1 transfected lysate using anti-SPINK1 monoclonal antibody and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with SPINK1 MaxPab rabbit polyclonal antibody.



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (31.9 KDa) .


Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



SPINK1 monoclonal antibody (M01), clone 4D4. Western Blot analysis of SPINK1 expression in human pancreas.



Immunoperoxidase of monoclonal antibody to SPINK1 on formalin-fixed paraffin-embedded human pancreas. [antibody concentration 1 ug/ml]



Detection limit for recombinant GST tagged SPINK1 is 0.03 ng/ml as a capture antibody.

SPINK1 Antibody (monoclonal) (M01) - Background

The protein encoded by this gene is a trypsin inhibitor, which is secreted from pancreatic acinar cells into pancreatic juice. It is thought

to function in the prevention of trypsin-catalyzed premature activation of zymogens within the pancreas and the pancreatic duct. Mutations in this gene are associated with hereditary pancreatitis and tropical calcific pancreatitis.

SPINK1 Antibody (monoclonal) (M01) - References

1.HOXB13 G84E-related Familial Prostate Cancers: A Clinical, Histologic, and Molecular Survey.Smith SC, Palanisamy N, Zuhlke KA, Johnson AM, Siddiqui J, Chinnaiyan AM, Kunju LP, Cooney KA, Tomlins SA. *Am J Surg Pathol*. 2014 May;38(5):615-26. doi: 10.1097/PAS.0000000000000090.

2.Novel dual-color immunohistochemical methods for detecting ERG-PTEN and ERG-SPINK1 status in prostate carcinoma.Bhalla R, Kunju LP, Tomlins SA, Christopherson K, Cortez C, Carskadon S, Siddiqui J, Park K, Miguel Mosquera J, Pestano GA, Rubin MA, Chinnaiyan AM, Palanisamy N. *Mod Pathol*. 2013 Jan 25. doi: 10.1038/modpathol.2012.234.

3.Loss of SPINK1 expression is associated with unfavorable outcomes in urothelial carcinoma of the bladder after radical cystectomy.Rink M, Park K, Volkmer BG, Xylinas E, Hansen J, Cha EK, Robinson BD, Hautmann R, Kufer R, Engel O, Chun FK, Dahlem R, Rubin MA, Shariat SF, Mosquera JM. *Urol Oncol*. 2012 Sep 1. pii: S1078-1439(12)00237-2. doi: 10.1016/j.urolonc.2012.06.011.

4.DIFFERENCES IN TMPRSS2-ERG GENE FUSION AND SPINK1 OVEREXPRESSION IN PROSTATE CANCER IN AFRICAN-AMERICAN AND CAUCASIAN MEN.Khani F, Mosquera JM, Park K, Srivastava A, Tewari A, Rubin M, Robinson B. *The Journal of Urology* Volume 187, Issue 4, Supplement, April 2012, Pages e1325.Characterization of ETS gene aberrations in select histologic variants of prostate carcinoma.Han B, Mehra R, Suleman K, Tomlins SA, Wang L, Singhal N, Linetzky KA, Palanisamy N, Zhou M, Chinnaiyan AM, Shah RB. *Mod Pathol*. 2009 Sep;22(9):1176-85. Epub 2009 May 22.

6.The role of SPINK1 in ETS rearrangement-negative prostate cancers.Tomlins SA, Rhodes DR, Yu J, Varambally S, Mehra R, Perner S, Demichelis F, Helgeson BE, Laxman B, Morris DS, Cao Q, Cao X, Andren O, Fall K, Johnson L, Wei JT, Shah RB, Al-Ahmadie H, Eastham JA, Egge SE, Fine SW, Hotakainen K, Stenman UH, Tsodikov

A,Cancer Cell. 2008 Jun;13(6):519-28.