

TIMP1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant TIMP1.

Catalog # AT4241a

Specification

TIMP1 Antibody (monoclonal) (M01) - Product Information

Application	IP, WB, E
Primary Accession	P01033
Other Accession	BC007097
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	23171

TIMP1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 7076

Other Names

Metalloproteinase inhibitor 1,
Erythroid-potentiating activity, EPA,
Fibroblast collagenase inhibitor,
Collagenase inhibitor, Tissue inhibitor of
metalloproteinases 1, TIMP-1, TIMP1, CLGI,
TIMP

Target/Specificity

TIMP1 (AAH07097, 1 a.a. ~ 169 a.a)
full-length recombinant protein with GST
tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format


Clear, colorless solution in phosphate
buffered saline, pH 7.2 .

Storage


Store at -20°C or lower. Aliquot to avoid
repeated freezing and thawing.

Precautions

TIMP1 Antibody (monoclonal) (M01) is for
research use only and not for use in
diagnostic or therapeutic procedures.



Immunoprecipitation of TIMP1 transfected lysate using anti-TIMP1 monoclonal antibody and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with TIMP1 MaxPab rabbit polyclonal antibody.




Western Blot analysis of TIMP1 expression in transfected 293T cell line by TIMP1 monoclonal antibody (M01), clone 4D12.

Lane 1: TIMP1 transfected lysate(23.2 KDa).
Lane 2: Non-transfected lysate.

TIMP1 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Detection limit for recombinant GST tagged TIMP1 is approximately 3ng/ml as a capture antibody.

TIMP1 Antibody (monoclonal) (M01) - Background

This gene belongs to the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix. In addition to its inhibitory role against most of the known MMPs, the encoded protein is able to promote cell proliferation in a wide range of cell types, and may also have an anti-apoptotic function. Transcription of this gene is highly inducible in response to many cytokines and hormones. In addition, the expression from some but not all inactive X chromosomes suggests that this gene inactivation is polymorphic in human females. This gene is located within intron 6 of the synapsin I gene and is transcribed in the opposite direction. [provided by RefSeq]