

TRIM23 Antibody (monoclonal) (M05)

Mouse monoclonal antibody raised against a partial recombinant TRIM23.

Catalog # AT4344a

Specification

TRIM23 Antibody (monoclonal) (M05) - Product Information

Application	WB, E
Primary Accession	P36406
Other Accession	BC022510
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	64067

TRIM23 Antibody (monoclonal) (M05) - Additional Information

Gene ID 373

Other Names

E3 ubiquitin-protein ligase TRIM23, 632-, ADP-ribosylation factor domain-containing protein 1, GTP-binding protein ARD-1, RING finger protein 46, Tripartite motif-containing protein 23, TRIM23, ARD1, ARFD1, RNF46

Target/Specificity

TRIM23 (AAH22510, 1 a.a. ~ 110 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format


Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage


Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

TRIM23 Antibody (monoclonal) (M05) is for research use only and not for use in diagnostic or therapeutic procedures.



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .



Detection limit for recombinant GST tagged TRIM23 is approximately 3ng/ml as a capture antibody.

TRIM23 Antibody (monoclonal) (M05) - Background

The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein is also a member of the ADP ribosylation factor family of guanine nucleotide-binding family of proteins. Its carboxy terminus contains an ADP-ribosylation factor domain and a guanine nucleotide binding site, while the amino

**TRIM23 Antibody (monoclonal) (M05) -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

terminus contains a GTPase activating protein domain which acts on the guanine nucleotide binding site. The protein localizes to lysosomes and the Golgi apparatus. It plays a role in the formation of intracellular transport vesicles, their movement from one compartment to another, and phospholipase D activation. Three alternatively spliced transcript variants for this gene have been described.

**TRIM23 Antibody (monoclonal) (M05) -
References**

Identification of TRIM23 as a cofactor involved in the regulation of NF-kappaB by human cytomegalovirus. Poole E, et al. J Virol, 2009 Apr. PMID 19176615. An empirical framework for binary interactome mapping. Venkatesan K, et al. Nat Methods, 2009 Jan. PMID 19060904. Towards a proteome-scale map of the human protein-protein interaction network. Rual JF, et al. Nature, 2005 Oct 20. PMID 16189514. E3 ubiquitin ligase activity of the trifunctional ARD1 (ADP-ribosylation factor domain protein 1). Vichi A, et al. Proc Natl Acad Sci U S A, 2005 Feb 8. PMID 15684077. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.