

Phospho-ErbB2(Y1112) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5151

Specification

Phospho-ErbB2(Y1112) Antibody - Product Information

Application	WB,E
Primary Accession	P04626
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=138 KDa
Isotype	Rabbit Ig
Antigen Source	HUMAN

Phospho-ErbB2(Y1112) Antibody - Additional Information

Gene ID 2064

Antigen Region
1100-1118

Other Names

Receptor tyrosine-protein kinase erbB-2 [Precursor]; p185erbB2; C-erbB-2; NEU proto-oncogene; Tyrosine kinase-type cell surface receptor HER2; MLN 19; ERBB2; HER2; NEU; NGL;

Dilution

WB~~1:500

Target/Specificity


This ErbB2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y1112 of human ErbB2.

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C



Western blot analysis of extracts from A431 cells, untreated or treated with EGF, 100ng/ml, using phospho-ErbB2 Antibody (Y1112) (upper) or Beta-actin (lower).



Western blot analysis of lysates from A431 cell line, untreated or treated with EGF, using phospho-ErbB2-Y1112 (Cat. #AW5151) (upper) or Beta-actin (lower).

Phospho-ErbB2(Y1112) Antibody -

in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-ErbB2(Y1112) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-ErbB2(Y1112) Antibody - Protein Information

Name ERBB2

Synonyms HER2, MLN19, NEU, NGL

Function

Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.

Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein Early endosome. Cytoplasm, perinuclear region. Nucleus. Note=Translocation to the nucleus requires endocytosis, probably endosomal sorting and is mediated by importin beta-1/KPNB1. Also detected in VPS35-positive endosome-to-TGN retrograde vesicles (PubMed:31138794). [Isoform 3]: Cytoplasm. Nucleus.

Tissue Location

Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.

Background

ErbB2 is a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors.

Phospho-ErbB2(Y1112) Antibody - References

Stephens, P., et al., Nature 431(7008):525-526 (2004). Wang, S.C., et al., Cancer Cell 6(3):251-261 (2004). Menendez, J.A., et al., Proc. Natl. Acad. Sci. U.S.A. 101(29):10715-10720 (2004). M, et al., Anticancer Res. 24(4):2219-2224 (2004). Contreras, D.N., et al., Clin. Appl. Thromb. Hemost. 10(3):271-276 (2004).

**Phospho-ErbB2(Y1112) Antibody -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)