

Phospho-SHP2(Y546) Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AW5158

Specification

Phospho-SHP2(Y546) Antibody - Product Information

Application WB, DB,E Primary Accession Q06124

Other Accession P41499, P35235,

Q90687

Reactivity Human, Mouse Predicted Chicken, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW H=68 KDa
Isotype Rabbit Ig
Antigen Source HUMAN

Phospho-SHP2(Y546) Antibody - Additional Information

Gene ID 5781

Antigen Region 521-551

Other Names

PTPN11; PTP2C; SHPTP2; Tyrosine-protein phosphatase non-receptor type 11; Protein-tyrosine phosphatase 1D; Protein-tyrosine phosphatase 2C; SH-PTP2; SH-PTP3

Dilution


WB~~1:1000 DB~~1:500

Target/Specificity


This SHP2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y546 of human SHP2.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.



Western blot analysis of lysate from NIH/3T3 cell line, using Phospho-SHP2-Y546 Antibody(Cat. #AW5158). AW5158 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.Lysate at 20ug.



Western blot analysis of lysates from NIH/3T3 cell line, untreated or treated with EGF, 100ng/ml, using Phospho-SHP2-Y546 Antibody(EV)(upper) or Beta-actin (lower).



Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-SHP2(Y546) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-SHP2(Y546) Antibody - Protein Information

Name PTPN11

Synonyms PTP2C, SHPTP2

Function

Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus (PubMed:10655584, PubMed:<a href="http://www.uniprot.org/citations/18559669"

target=" blank">18559669,

PubMed: <a href="http://www.uniprot.org/ci tations/18829466"

target="_blank">18829466,

PubMed:<a href="http://www.uniprot.org/ci tations/26742426"

target=" blank">26742426,

PubMed:<a href="http://www.uniprot.org/ci tations/28074573"

target=" blank">28074573).

Positively regulates MAPK signal

transduction pathway (PubMed:<a href="ht tp://www.uniprot.org/citations/28074573"

target=" blank">28074573).

Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed:http://www.uniprot

.org/citations/28074573"

target="_blank">28074573).

Dephosphorylates ROCK2 at 'Tyr-722'

resulting in stimulation of its RhoA binding activity (PubMed:<a href="http://www.unipr

ot.org/citations/18559669"


target="_blank">18559669).

Dephosphorylates CDC73 (PubMed:<a href = "http://www.uniprot.org/citations/2674242"

6" target="_blank">26742426).

Dephosphorylates SOX9 on tyrosine

residues, leading to inactivate SOX9 and promote ossification (By similarity).



Dot blot analysis of anti-Phospho-SHP2-Y546 Phospho-specific Pab (Cat. #AW5158) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.


Phospho-SHP2(Y546) Antibody - Background

SHP2 is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains two tandem Src homology-2 domains, which function as phospho-tyrosine binding domains and mediate the interaction of this PTP with its substrates. This PTP is widely expressed in most tissues and plays a regulatory role in various cell signaling events that are important for a diversity of cell functions, such as mitogenic activation, metabolic control, transcription regulation, and cell migration.

Phospho-SHP2(Y546) Antibody - References

Nystrom,A.M., et.al., Eur J Med Genet (2010) In press

Carver, K.C., et.al., J. Biol. Chem. 285 (11), 8003-8012 (2010)





Cellular Location Cytoplasm. Nucleus

Tissue LocationWidely expressed, with highest levels in heart, brain, and skeletal muscle.

Phospho-SHP2(Y546) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture