

UCHL1 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5247

Specification

UCHL1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P09936
Other Accession	Q00981 , Q6SEG5 , Q9R0P9 , Q60HC8 , P23356 , Q9GM50
Reactivity	Human, Mouse, Rat
Predicted	Bovine, Horse, Monkey, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=25;M=25;Rat= 25 KDa
Isotype	Rabbit Ig
Antigen Source	HUMAN

UCHL1 Antibody (C-term) - Additional Information

Gene ID 7345

Antigen Region
187-216

Other Names

UCHL1; Ubiquitin carboxyl-terminal hydrolase isozyme L1; Neuron cytoplasmic protein 9.5; PGP 9.5; Ubiquitin thioesterase L1

Dilution


WB~~1:1000

Target/Specificity


This UCHL1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 187-216 amino acids from the C-terminal region of human UCHL1.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.



Western blot analysis of lysates from mouse brain tissue, rat PC-12, rat C6, mouse Neuro-2a, SH-SY5Y cell line (from left to right), using UCHL1 Antibody (C-term) (Cat. #AW5247). AW5247 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody.



Western blot analysis of lysates from mouse Neuro-2a cell line, mouse brain tissue lysate, rat PC-12, rat C6 cell line (from left to

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UCHL1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

UCHL1 Antibody (C-term) - Protein Information**Name** UCHL1**Function**

Ubiquitin-protein hydrolase involved both in the processing of ubiquitin precursors and of ubiquitinated proteins (Probable). This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin (PubMed:9774100, PubMed:8639624, PubMed:12408865, PubMed:23359680). Also binds to free monoubiquitin and may prevent its degradation in lysosomes (By similarity). The homodimer may have ATP-independent ubiquitin ligase activity (PubMed:12408865).

Cellular Location

Cytoplasm. Endoplasmic reticulum membrane; Lipid- anchor. Note=About 30% of total UCHL1 is associated with membranes in brain

Tissue Location

Found in neuronal cell bodies and processes throughout the neocortex (at protein level). Expressed in neurons and cells of the diffuse neuroendocrine system and their tumors. Weakly expressed in ovary. Down-regulated in brains from Parkinson disease and Alzheimer disease patients.

right), using UCHL1 Antibody (C-term)(Cat. #AW5247). AW5247 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.

UCHL1 Antibody (C-term) - Background

UCHL1 is a member of a gene family whose products hydrolyze small C-terminal adducts of ubiquitin to generate the ubiquitin monomer. Expression of UCHL1 is highly specific to neurons and to cells of the diffuse neuroendocrine system and their tumors. It is present in all neurons (Doran et al., 1983 [PubMed 6343558]).

UCHL1 Antibody (C-term) - References

Maraganore, D.M., et al., Mov Disord 18(6):631-636 (2003). Nishikawa, K., et al., Biochem. Biophys. Res. Commun. 304(1):176-183 (2003). Liu, Y., et al., Cell 111(2):209-218 (2002). Caballero, O.L., et al., Oncogene 21(19):3003-3010 (2002). Saigoh, K., et al., Nat. Genet. 23(1):47-51 (1999).

UCHL1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)