

ERBB2 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AW5248

Specification

ERBB2 Antibody - Product Information

Application	WB,E
Primary Accession	P04626
Reactivity	Human
Host	Mouse
Clonality	monoclonal
Calculated MW	H=140,71,63,136, 135,97 KDa
Isotype	IgG2b,κ
Antigen Source	HUMAN

ERBB2 Antibody - Additional Information

Gene ID 2064

Other Names

Receptor tyrosine-protein kinase erbB-2,
Metastatic lymph node gene 19 protein,
MLN 19, Proto-oncogene Neu,
Proto-oncogene c-ErbB-2, Tyrosine
kinase-type cell surface receptor HER2,
p185erbB2, CD340, ERBB2, HER2, MLN19,
NEU, NGL

Dilution

WB~~1:1000

Target/Specificity

This ERBB2 antibody is generated from a mouse immunized with a recombinant protein.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ERBB2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

ERBB2 Antibody - Protein Information



Western blot analysis of lysate from SK-BR-3,MDA-MB-453 cell line (from left to right), using ERBB2 Antibody (Cat. #AW5248). AW5248 was diluted at 1:1000. A goat anti-mouse IgG H&L(HRP) at 1:5,000 dilution was used as the secondary antibody. Lysate at 20ug per lane.

ERBB2 Antibody - Background

Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.

Name ERBB2**Synonyms** HER2, MLN19, NEU, NGL**Function**

Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.

Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein Early endosome. Cytoplasm, perinuclear region. Nucleus. Note=Translocation to the nucleus requires endocytosis, probably endosomal sorting and is mediated by importin beta-1/KPNB1. Also detected in VPS35-positive endosome-to-TGN retrograde vesicles (PubMed:31138794). [Isoform 3]: Cytoplasm. Nucleus.

Tissue Location

Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.

ERBB2 Antibody - References

Yamamoto T.,et al.Nature 319:230-234(1986).
Coussens L.,et al.Science 230:1132-1139(1985).
Wakamatsu A.,et al.Submitted (OCT-2007) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Tal M.,et al.Mol. Cell. Biol. 7:2597-2601(1987).

ERBB2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)