

(Mouse) Shh Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5359

Specification

(Mouse) Shh Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q62226
Reactivity	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	M=48;Rat=48
	KDa
Isotype	Rabbit Ig
Antigen Source	HUMAN

(Mouse) Shh Antibody (C-term) - Additional Information

Gene ID 20423

Antigen Region
397-431

Other Names

Sonic hedgehog protein, SHH, HHG-1, Sonic hedgehog protein N-product, Sonic hedgehog protein C-product, Shh, Hhg1

Dilution

WB~~1:1000
IHC-P~~1:25

Target/Specificity


This (Mouse) Shh antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 397-431 amino acids from the C-terminal region of Mouse Shh.

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C



All lanes : Anti-Shh Antibody (C-term)(AW5359) at 1/1000 dilution Lane 1: mouse liver lysates Lane 2: mouse lung lysates Lane 3: mouse stomach lysates Lane 4: rat lung lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 57 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded M. lung section using

in small aliquots to prevent freeze-thaw cycles.

Precautions

(Mouse) Shh Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

(Mouse) Shh Antibody (C-term) - Protein Information

Name Shh {ECO:0000312|MGI:MGI:98297}

Synonyms Hhgl1

Function

[Sonic hedgehog protein]: The C-terminal part of the sonic hedgehog protein precursor displays an autoproteolysis and a cholesterol transferase activity (PubMed:8824192, PubMed:7891723). Both activities result in the cleavage of the full-length protein into two parts (ShhN and ShhC) followed by the covalent attachment of a cholesterol moiety to the C-terminal of the newly generated ShhN (PubMed:8824192). Both activities occur in the reticulum endoplasmic (PubMed:21357747). Once cleaved, ShhC is degraded in the endoplasmic reticulum (PubMed:21357747).

Cellular Location

[Sonic hedgehog protein N-product]: Cell membrane; Lipid-anchor. Note=The dual-lipidated sonic hedgehog protein N-product (ShhNp) is firmly tethered to the cell membrane where it forms multimers (PubMed:24522195). Further solubilization and release from the cell surface seem to be achieved through different mechanisms, including the interaction with DISP1 and SCUBE2, movement by lipoprotein particles, transport by cellular extensions called cytonemes or by the proteolytic removal of both terminal lipidated peptides.

(Mouse) Shh Antibody (C-term)(Cat#AW5359). AW5359 was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

(Mouse) Shh Antibody (C-term) - Background

Binds to the patched (PTC) receptor, which functions in association with smoothened (SMO), to activate the transcription of target genes. In the absence of SHH, PTC represses the constitutive signaling activity of SMO. Also regulates another target, the gli oncogene. Intercellular signal essential for a variety of patterning events during development: signal produced by the notochord that induces ventral cell fate in the neural tube and somites, and the polarizing signal for patterning of the anterior-posterior axis of the developing limb bud. Displays both floor plate- and motor neuron-inducing activity. The threshold concentration of N-product required for motor neuron induction is 5-fold lower than that required for floor plate induction (By similarity).

(Mouse) Shh Antibody (C-term) - References

Echelard Y.,et al.Cell 75:1417-1430(1993).
McMahon A.P.,et al.Submitted (NOV-1997) to the EMBL/GenBank/DDBJ databases.
Chang D.T.,et al.Development 120:3339-3353(1994).
Carninci P.,et al.Science 309:1559-1563(2005).
Roelink H.,et al.Cell 81:445-455(1995).

Tissue Location

Expressed in a number of embryonic tissues including the notochord, ventral neural tube, floor plate, lung bud, zone of polarizing activity and posterior distal mesenchyme of limbs In the adult, expressed in lung and neural retina

(Mouse) Shh Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)