

NCAM1 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5412

Specification

NCAM1 Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P13591
Other Accession	P13596 , P13595 , P13590 , P31836
Reactivity Predicted	Human, Rat Bovine, Chicken, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=95,93;M=119,9 4;R=95 KDa
Isotype	Rabbit Ig
Antigen Source	HUMAN

NCAM1 Antibody (C-term) - Additional Information

Gene ID 4684


Antigen Region
796-830

Other Names
Neural cell adhesion molecule 1, N-CAM-1,
NCAM-1, CD56, NCAM1, NCAM


Dilution
WB~~1:1000
IHC-P~~1:25
FC~~1:25

Target/Specificity
This NCAM1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 796-830 amino acids from the C-terminal region of human NCAM1.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.



All lanes : Anti-NCAM1 Antibody (C-term) at 1:1000 dilution Lane 1: HT-1080 whole cell lysates Lane 2: SW480 whole cell lysates Lane 3: human cerebellum lysates Lane 4: human brain lysates Lane 5: rat brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 95 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Immunohistochemical analysis of

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NCAM1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NCAM1 Antibody (C-term) - Protein Information

Name NCAM1

Synonyms NCAM

Function

This protein is a cell adhesion molecule involved in neuron- neuron adhesion, neurite fasciculation, outgrowth of neurites, etc. (Microbial infection) Acts as a receptor for Zika virus.

Cellular Location


[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Isoform 3]: Cell membrane; Lipid-anchor, GPI- anchor [Isoform 5]: Secreted.

NCAM1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

paraffin-embedded H. astrogloma section using NCAM1 Antibody (C-term)(Cat#AW5412). AW5412 was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Flow cytometric analysis of SH-SY5Y cells using NCAM1 Antibody (C-term)(green, Cat#AW5412) compared to an isotype control of rabbit IgG(blue). AW5412 was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

NCAM1 Antibody (C-term) - Background

This protein is a cell adhesion molecule involved in neuron-neuron adhesion, neurite fasciculation, outgrowth of neurites, etc.

NCAM1 Antibody (C-term) - References

Barton C.H.,et al.Development 104:165-173(1988).
Lanier L.L.,et al.J. Immunol. 146:4421-4426(1991).
Saito S.,et al.Lung Cancer 10:307-318(1994).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Totoki Y.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.