

MAOA Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5425

Specification

MAOA Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P21397
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=60,45;M=60;R=60 KDa
Isotype	Rabbit Ig
Antigen Source	HUMAN

MAOA Antibody (C-term) - Additional Information

Gene ID 4128

Antigen Region
465-499

Other Names
Amine oxidase [flavin-containing] A,
Monoamine oxidase type A, MAO-A, MAOA

Dilution
WB~~1:1000

Target/Specificity


This MAOA antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 465-499 amino acids from the C-terminal region of human MAOA.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



All lanes : Anti-MAOA Antibody (C-term) at 1:1000 dilution Lane 1: human placenta lysates Lane 2: HepG2 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 60 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

MAOA Antibody (C-term) - Background

Catalyzes the oxidative deamination of biogenic and xenobiotic amines and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues. MAOA preferentially oxidizes biogenic amines such as 5-hydroxytryptamine (5-HT), norepinephrine and epinephrine.

MAOA Antibody (C-term) - References

Hsu Y.-P.P.,et al.J. Neurochem.
51:1321-1324(1988).
Bach A.W.J.,et al.Proc. Natl. Acad. Sci. U.S.A.
85:4934-4938(1988).
Chen Z.-Y.,et al.Nucleic Acids Res.
19:4537-4541(1991).

Precautions

MAOA Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Grimsby J., et al. Proc. Natl. Acad. Sci. U.S.A. 88:3637-3641(1991).

Ota T., et al. Nat. Genet. 36:40-45(2004).

MAOA Antibody (C-term) - Protein Information**Name** MAOA**Function**

Catalyzes the oxidative deamination of biogenic and xenobiotic amines and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues. MAOA preferentially oxidizes biogenic amines such as 5-hydroxytryptamine (5-HT), norepinephrine and epinephrine.

Cellular Location

Mitochondrion outer membrane; Single-pass type IV membrane protein; Cytoplasmic side

Tissue Location

Heart, liver, duodenum, blood vessels and kidney.

MAOA Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)