

SNAP25 Antibody (S187)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5468

Specification

SNAP25 Antibody (S187) - Product Information

Application	WB,E
Primary Accession	P60880
Other Accession	P60881 , P60879 , P60878 , Q170Q3 , Q5TZ66
Reactivity Predicted	Human, Mouse Zebrafish, Bovine, Chicken, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=23;M=23;R=23 KDa
Isotype	Rabbit Ig
Antigen Source	HUMAN

SNAP25 Antibody (S187) - Additional Information

Gene ID 6616

Antigen Region
179-213

Other Names

Synaptosomal-associated protein 25,
SNAP-25, Super protein, SUP,
Synaptosomal-associated 25 kDa protein,
SNAP25, SNAP

Dilution

WB~~1:1000


Target/Specificity

This SNAP25 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 179-213 amino acids from the region of human SNAP25.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage



All lanes : Anti-SNAP25 Antibody (S187) at 1:1000 dilution Lane 1: mouse brain lysates Lane 2: mouse cerebellum lysates Lane 3: human brain lysates Lane 4: human cerebellum lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 23 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

SNAP25 Antibody (S187) - Background

t-SNARE involved in the molecular regulation of neurotransmitter release. May play an important role in the synaptic function of specific neuronal systems. Associates with proteins involved in vesicle docking and membrane fusion. Regulates plasma membrane recycling through its interaction with CENPF.

SNAP25 Antibody (S187) - References

Bark I.C.,et al.Gene 139:291-292(1994).
Zhao N.,et al.Gene 145:313-314(1994).
Jagadish M.N.,et al.Biochem. J. 317:945-954(1996).
Kalnine N.,et al.Submitted (OCT-2004) to the

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).

Precautions

SNAP25 Antibody (S187) is for research use only and not for use in diagnostic or therapeutic procedures.

SNAP25 Antibody (S187) - Protein Information

Name SNAP25

Synonyms SNAP

Function

t-SNARE involved in the molecular regulation of neurotransmitter release. May play an important role in the synaptic function of specific neuronal systems. Associates with proteins involved in vesicle docking and membrane fusion. Regulates plasma membrane recycling through its interaction with CENPF. Modulates the gating characteristics of the delayed rectifier voltage-dependent potassium channel KCNB1 in pancreatic beta cells.

Cellular Location

Cytoplasm, perinuclear region
{ECO:0000250|UniProtKB:P60879}. Cell membrane
{ECO:0000250|UniProtKB:P60881};
Lipid-anchor
{ECO:0000250|UniProtKB:P60879}. Cell junction, synapse, synaptosome
{ECO:0000250|UniProtKB:P60879}. Photoreceptor inner segment
{ECO:0000250|UniProtKB:P60879}.
Note=Membrane association requires palmitoylation. Expressed throughout cytoplasm, concentrating at the perinuclear region. Colocalizes with KCNB1 at the cell membrane (By similarity). Colocalizes with PLCL1 at the cell membrane (By similarity).
{ECO:0000250|UniProtKB:P60879, ECO:0000250|UniProtKB:P60881}

Tissue Location

Neurons of the neocortex, hippocampus, piriform cortex, anterior thalamic nuclei, pontine nuclei, and granule cells of the cerebellum

SNAP25 Antibody (S187) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)