

**TMED10 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
Catalog # AW5552

**Specification**

**TMED10 Antibody (C-term) - Product Information**

Application	<b>WB,E</b>
Primary Accession	<a href="#">P49755</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>H=25;M=25,11;R=25(20kd□□) KDa</b>
Isotype	<b>Rabbit Ig</b>
Antigen Source	<b>HUMAN</b>

**TMED10 Antibody (C-term) - Additional Information**

**Gene ID** 10972

**Antigen Region**  
156-185

**Other Names**

Transmembrane emp24 domain-containing protein 10, 21 kDa  
transmembrane-trafficking protein, S31III125, S31I125, Tmp-21-I,  
Transmembrane protein Tmp21, p23, p24 family protein delta-1, p24delta1, p24delta, TMED10, TMP21

**Dilution**

WB~~1:2000


**Target/Specificity**

This TMED10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 156-185 amino acids from the C-terminal region of human TMED10.


**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**



All lanes : Anti-TMED10 Antibody (C-term) at 1:1000 dilution Lane 1: PC-12 whole cell lysate Lane 2: NCI-H460 whole cell lysate Lane 3: HeLa whole cell lysate Lane 4: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes : Anti-TMED10 Antibody (C-term) at 1:1000 dilution Lane 1: NIH/3T3 whole cell

TMED10 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### TMED10 Antibody (C-term) - Protein Information


Name TMED10 ([HGNC:16998](#))

Synonyms TMP21

#### Function

Cargo receptor involved in protein vesicular trafficking and quality control in the endoplasmic reticulum (ER) and Golgi (PubMed: [10052452](http://www.uniprot.org/citations/10052452)) target="\_blank">10052452</a>, PubMed: [11726511](http://www.uniprot.org/citations/11726511) target="\_blank">11726511</a>, PubMed: [16641999](http://www.uniprot.org/citations/16641999) target="\_blank">16641999</a>, PubMed: [17288597](http://www.uniprot.org/citations/17288597) target="\_blank">17288597</a>, PubMed: [19296914](http://www.uniprot.org/citations/19296914) target="\_blank">19296914</a>, PubMed: [20427317](http://www.uniprot.org/citations/20427317) target="\_blank">20427317</a>, PubMed: [21219331](http://www.uniprot.org/citations/21219331) target="\_blank">21219331</a>, PubMed: [27569046](http://www.uniprot.org/citations/27569046) target="\_blank">27569046</a>). The p24 protein family is a group of transmembrane proteins that bind coat protein complex I/COPI and coat protein complex II/COPII involved in vesicular trafficking between the membranes (PubMed: [10052452](http://www.uniprot.org/citations/10052452)) target="\_blank">10052452</a>). Acts at the luminal side for incorporation of secretory cargo molecules into transport vesicles and involved in vesicle coat formation at the cytoplasmic side (PubMed: [20427317](http://www.uniprot.org/citations/20427317)) target="\_blank">20427317</a>, PubMed: [27569046](http://www.uniprot.org/citations/27569046) target="\_blank">27569046</a>). Mainly functions in the early secretory pathway

lysate Lane 2: PC-12 whole cell lysate Lane 3: NCI-H460 whole cell lysate Lane 4: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-TMED10 Antibody (C-term) at 1:2000 dilution Lane 1: human pancreas lysate Lane 2: HeLa whole cell lysate Lane 3: NCI-H460 whole cell lysate Lane 4: Neuro-2a whole cell lysate Lane 5: NIH/3T3 whole cell lysate Lane 5: PC-12 whole cell lysate Lane 5: U-87 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

#### TMED10 Antibody (C-term) - Background

This gene is a member of the EMP24/GP25L/p24 family and encodes a protein with a GOLD domain. This type I membrane protein is localized to the plasma membrane and golgi cisternae and is involved in vesicular protein trafficking. The protein is also a member of a heteromeric secretase complex and regulates the complex's gamma-secretase activity without affecting its epsilon-secretase activity. Mutations in this gene have been associated with early-onset familial Alzheimer's disease. This gene

and cycles between the ER, ER-Golgi intermediate compartment (ERGIC) and Golgi, mediating cargo transport through COPI and COPII-coated vesicles (PubMed:<a href="http://www.uniprot.org/citations/10052452" target="\_blank">10052452</a>, PubMed:<a href="http://www.uniprot.org/citations/10852829" target="\_blank">10852829</a>, PubMed:<a href="http://www.uniprot.org/citations/12237308" target="\_blank">12237308</a>). In COPII vesicle-mediated anterograde transport, involved in the transport of GPI-anchored proteins by acting together with TMED2 as their cargo receptor; the function specifically implies SEC24C and SEC24D of the COPII vesicle coat and lipid raft-like microdomains of the ER (PubMed:<a href="http://www.uniprot.org/citations/20427317" target="\_blank">20427317</a>, PubMed:<a href="http://www.uniprot.org/citations/27569046" target="\_blank">27569046</a>). Recognizes GPI anchors structural remodeled in the ER by the GPI inositol-deacylase/PGAP1 and the metallophosphoesterase MPPE1/PGAP5 (By similarity). In COPI vesicle-mediated retrograde transport, involved in the biogenesis of COPI vesicles and vesicle coat recruitment (PubMed:<a href="http://www.uniprot.org/citations/11726511" target="\_blank">11726511</a>). Involved in trafficking of amyloid beta A4 protein and soluble APP-beta release (independent from the modulation of gamma-secretase activity) (PubMed:<a href="http://www.uniprot.org/citations/17288597" target="\_blank">17288597</a>). Involved in the KDEL2-mediated retrograde transport of the toxin A subunit (CTX-A-K63) together with COPI and the COOH terminus of KDEL2 (By similarity). On Golgi membranes, acts as primary receptor for ARF1-GDP, a GTP-binding protein involved in COPI-vesicle formation (PubMed:<a href="http://www.uniprot.org/citations/11726511" target="\_blank">11726511</a>). Increases coatomer-dependent GTPase-activating activity of ARFGAP2 which mediates the hydrolysis of ARF1-bound GTP and therefore modulates protein trafficking from the Golgi apparatus (PubMed:<a href="http://www.uniprot.org/citations/19296914" target="\_blank">19296914</a>). Involved

has a pseudogene on chromosome 8.

#### TMED10 Antibody (C-term) - References

Wang, H., et al. Mol. Biol. Cell 21(8):1398-1408(2010)  
Zhao, J., et al. BMC Med. Genet. 11, 96 (2010) :  
Pardossi-Piquard, R., et al. J. Biol. Chem. 284(42):28634-28641(2009)  
Soranzo, N., et al. PLoS Genet. 5 (4), E1000445 (2009) :  
Liu, S., et al. Eur. J. Neurosci. 28(10):1980-1988(2008)

in the exocytic trafficking of G protein-coupled receptors F2LR1/PAR2 (trypsin and trypsin-like enzyme receptor), OPRM1 (opioid receptor) and P2RY4 (UTD and UDP receptor) from the Golgi to the plasma membrane, thus contributing to receptor resensitization (PubMed:<a href="http://www.uniprot.org/citations/21219331" target="\_blank">21219331</a>). In addition to its cargo receptor activity, may also act as a protein channel after oligomerization, facilitating the post-translational entry of leaderless cytoplasmic cargo into the ERGIC (PubMed:<a href="http://www.uniprot.org/citations/32272059" target="\_blank">32272059</a>). Involved in the translocation into ERGIC, the vesicle entry and the secretion of leaderless cargos (lacking the secretion signal sequence), including the mature form of interleukin 1/IL-1 family members, the alpha-crystallin B chain HSPB5, the carbohydrate-binding proteins galectin-1/LGALS1 and galectin-3/LGALS3, the microtubule-associated protein Tau/MAPT, and the annexin A1/ANXA1; the translocation process is dependent on cargo protein unfolding and enhanced by chaperones HSP90AB1 and HSP90B1/GRP9 (PubMed:<a href="http://www.uniprot.org/citations/32272059" target="\_blank">32272059</a>). Could also associates with the presenilin-dependent gamma-secretase complex in order to regulate gamma-cleavages of the amyloid beta A4 protein to yield amyloid-beta 40/Abeta40 (PubMed:<a href="http://www.uniprot.org/citations/16641999" target="\_blank">16641999</a>).

#### Cellular Location

Endoplasmic reticulum membrane; Single-pass type I membrane protein. Endoplasmic reticulum-Golgi intermediate compartment membrane; Single-pass type I membrane protein. Golgi apparatus membrane; Single-pass type I membrane protein. Golgi apparatus, cis-Golgi network membrane; Single-pass type I membrane protein. Golgi apparatus, trans-Golgi network membrane {ECO:0000250|UniProtKB:Q63584}; Single-pass type I membrane protein. Cytoplasmic vesicle, secretory vesicle membrane; Single-pass type I membrane protein. Cell membrane

{ECO:0000250|UniProtKB:Q63584};  
Single-pass type I membrane protein.  
Melanosome Note=Identified by mass  
spectrometry in melanosome fractions from  
stage I to stage IV.

### **TMED10 Antibody (C-term) - Protocols**

Provided below are standard protocols that you  
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)