

TRIM23 Antibody (N-term) Blocking peptide
Synthetic peptide
Catalog # BP13641a**Specification****TRIM23 Antibody (N-term) Blocking peptide -
Product Information**Primary Accession [P36406](#)**TRIM23 Antibody (N-term) Blocking peptide -
Additional Information****Gene ID 373****Other Names**

E3 ubiquitin-protein ligase TRIM23, 632-, ADP-ribosylation factor domain-containing protein 1, GTP-binding protein ARD-1, RING finger protein 46, Tripartite motif-containing protein 23, TRIM23, ARD1, ARFD1, RNF46

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13641a was selected from the N-term region of TRIM23. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**TRIM23 Antibody (N-term) Blocking peptide -
Protein Information****Name** TRIM23**TRIM23 Antibody (N-term) Blocking
peptide - Background**

The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein is also a member of the ADP-ribosylation factor family of guanine nucleotide-binding family of proteins. Its carboxy terminus contains an ADP-ribosylation factor domain and a guanine nucleotide binding site, while the amino terminus contains a GTPase activating protein domain which acts on the guanine nucleotide binding site. The protein localizes to lysosomes and the Golgi apparatus. It plays a role in the formation of intracellular transport vesicles, their movement from one compartment to another, and phospholipase D activation. Three alternatively spliced transcript variants for this gene have been described.

**TRIM23 Antibody (N-term) Blocking
peptide - References**

Arimoto, K., et al. Proc. Natl. Acad. Sci. U.S.A. 107(36):15856-15861(2010) Poole, E., et al. J. Virol. 83(8):3581-3590(2009) Venkatesan, K., et al. Nat. Methods 6(1):83-90(2009) Vichi, A., et al. Proc. Natl. Acad. Sci. U.S.A. 102(6):1945-1950(2005) Raymond, A., et al. EMBO J. 20(9):2140-2151(2001)

Synonyms ARD1, ARFD1, RNF46

Function

Acts as an E3 ubiquitin-protein ligase. Plays an essential role in autophagy activation during viral infection. Mechanistically, activates TANK-binding kinase 1/TBK1 by facilitating its dimerization and ability to phosphorylate the selective autophagy receptor SQSTM1. In order to achieve this function, TRIM23 mediates 'Lys-27'-linked auto-ubiquitination of its ADP-ribosylation factor (ARF) domain to induce its GTPase activity and its recruitment to autophagosomes (PubMed:28871090).

Cellular Location

Cytoplasm. Endomembrane system. Golgi apparatus membrane. Lysosome membrane. Note=Membrane-associated with the Golgi complex and lysosomal structures

TRIM23 Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)