

PTEN Antibody (C-term S362) Blocking Peptide
Synthetic peptide
Catalog # BP18873b**Specification****PTEN Antibody (C-term S362) Blocking Peptide - Product Information**Primary Accession [P60484](#)**PTEN Antibody (C-term S362) Blocking Peptide - Additional Information****Gene ID** 5728**Other Names**

Phosphatidylinositol 3, 5-trisphosphate 3-phosphatase and dual-specificity protein phosphatase PTEN, Mutated in multiple advanced cancers 1, Phosphatase and tensin homolog, PTEN, MMAC1, TEP1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PTEN Antibody (C-term S362) Blocking Peptide - Protein Information**Name** PTEN**Synonyms** MMAC1, TEP1**Function**

Tumor suppressor. Acts as a dual-specificity protein phosphatase, dephosphorylating tyrosine-, serine- and threonine-phosphorylated proteins. Also acts as a lipid phosphatase, removing the phosphate in

PTEN Antibody (C-term S362) Blocking Peptide - Background

This gene was identified as a tumor suppressor that is mutated in a large number of cancers at high frequency. The protein encoded by this gene is a phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase. It contains a tensin-like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. Unlike most of the protein tyrosine phosphatases, this protein preferentially dephosphorylates phosphoinositide substrates. It negatively regulates intracellular levels of phosphatidylinositol-3,4,5-trisphosphate in cells and functions as a tumor suppressor by negatively regulating AKT/PKB signaling pathway.

PTEN Antibody (C-term S362) Blocking Peptide - References

Miletic, A.V., et al. J. Exp. Med. 207(11):2407-2420(2010)Kini, V., et al. J. Biol. Chem. 285(43):33082-33091(2010)Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)Molina, J.R., et al. Cancer Res. 70(17):6697-6703(2010)Iliopoulos, D., et al. Mol. Cell 39(4):493-506(2010)

the D3 position of the inositol ring from phosphatidylinositol 3,4,5-trisphosphate, phosphatidylinositol 3,4- diphosphate, phosphatidylinositol 3-phosphate and inositol 1,3,4,5- tetrakisphosphate with order of substrate preference in vitro $\text{PtdIns}(3,4,5)\text{P}_3 > \text{PtdIns}(3,4)\text{P}_2 > \text{PtdIns}3\text{P} > \text{Ins}(1,3,4,5)\text{P}_4$ (PubMed:26504226, PubMed:16824732). The lipid phosphatase activity is critical for its tumor suppressor function. Antagonizes the PI3K-AKT/PKB signaling pathway by dephosphorylating phosphoinositides and thereby modulating cell cycle progression and cell survival. The unphosphorylated form cooperates with MAGI2 to suppress AKT1 activation. Dephosphorylates tyrosine-phosphorylated focal adhesion kinase and inhibits cell migration and integrin-mediated cell spreading and focal adhesion formation. Plays a role as a key modulator of the AKT-mTOR signaling pathway controlling the tempo of the process of newborn neurons integration during adult neurogenesis, including correct neuron positioning, dendritic development and synapse formation. May be a negative regulator of insulin signaling and glucose metabolism in adipose tissue. The nuclear monoubiquitinated form possesses greater apoptotic potential, whereas the cytoplasmic nonubiquitinated form induces less tumor suppressive ability. In motile cells, suppresses the formation of lateral pseudopods and thereby promotes cell polarization and directed movement.

Cellular Location

Cytoplasm. Nucleus. Nucleus, PML body.
Note=Monoubiquitinated form is nuclear.
Nonubiquitinated form is cytoplasmic.
Colocalized with PML and USP7 in PML nuclear bodies (PubMed:18716620).
XIAP/BIRC4 promotes its nuclear localization (PubMed:19473982).

Tissue Location

Expressed at a relatively high level in all adult tissues, including heart, brain, placenta, lung, liver, muscle, kidney and pancreas.

PTEN Antibody (C-term S362) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)