

IKKB Antibody (C-term S692) Blocking Peptide
Synthetic peptide
Catalog # BP18876b**Specification****IKKB Antibody (C-term S692) Blocking Peptide - Product Information**Primary Accession [O14920](#)**IKKB Antibody (C-term S692) Blocking Peptide - Additional Information****Gene ID** 3551**Other Names**

Inhibitor of nuclear factor kappa-B kinase subunit beta, I-kappa-B-kinase beta, IKK-B, IKK-beta, IkbKB, I-kappa-B kinase 2, IKK2, Nuclear factor NF-kappa-B inhibitor kinase beta, NFKBIKB, IKBKB, IKKB

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

IKKB Antibody (C-term S692) Blocking Peptide - Protein Information**Name** IKBKB**Synonyms** IKKB**Function**

Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular

IKKB Antibody (C-term S692) Blocking Peptide - Background

NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL(MIM 164910), RELA (MIM 164014), or RELB (MIM 604758) to form theNFKB complex. The NFKB complex is inhibited by I-kappa-B proteins(NFKBIA, MIM 164008, or NFKBIB, MIM 604495), which inactivateNF-kappa-B by trapping it in the cytoplasm. Phosphorylation ofserine residues on the I-kappa-B proteins by kinases (IKBKA, MIM600664, or IKBKB) marks them for destruction via the ubiquitinationpathway, thereby allowing activation of the NF-kappa-B complex.Activated NFKB complex translocates into the nucleus and binds DNAat kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or Gpurine; and Y is a C or T pyrimidine).

IKKB Antibody (C-term S692) Blocking Peptide - References

Yatherajam, G., et al. J. Immunol. 185(5):2665-2669(2010)Kenneth, N.S., et al. EMBO J. 29(17):2966-2978(2010)Zhao, M., et al. J. Biol. Chem. 285(32):24372-24380(2010)Niida, M., et al. Mol. Immunol. 47(14):2378-2387(2010)Schoorhof, A., et al. Pediatr. Pulmonol. 45(6):608-613(2010)

stresses (PubMed:30337470). Acts as part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation. Phosphorylates inhibitors of NF-kappa-B on 2 critical serine residues. These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome. In turn, free NF-kappa-B is translocated into the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis. In addition to the NF-kappa-B inhibitors, phosphorylates several other components of the signaling pathway including NEMO/IKBKG, NF-kappa-B subunits RELA and NFkB1, as well as IKK-related kinases TBK1 and IKBKE (PubMed:11297557, PubMed:20410276). IKK-related kinase phosphorylations may prevent the overproduction of inflammatory mediators since they exert a negative regulation on canonical IKKs. Phosphorylates FOXO3, mediating the TNF-dependent inactivation of this pro-apoptotic transcription factor (PubMed:15084260). Also phosphorylates other substrates including NCOA3, BCL10 and IRS1 (PubMed:17213322). Within the nucleus, acts as an adapter protein for NFkBIA degradation in UV-induced NF-kappa-B activation (PubMed:11297557). Phosphorylates RIPK1 at 'Ser-25' which represses its kinase activity and consequently prevents TNF-mediated RIPK1-dependent cell death (By similarity). Phosphorylates the C- terminus of IRF5, stimulating IRF5 homodimerization and translocation into the nucleus (PubMed:25326418).

Cellular Location

Cytoplasm. Nucleus. Membrane raft.

Note=Colocalized with DPP4 in membrane

rafts.

Tissue Location

Highly expressed in heart, placenta, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis and peripheral blood

IKKB Antibody (C-term S692) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)