

**CHST2 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP19294b****Specification****CHST2 Antibody (C-term) Blocking Peptide -  
Product Information**Primary Accession [Q9Y4C5](#)**CHST2 Antibody (C-term) Blocking Peptide -  
Additional Information****Gene ID** 9435**Other Names**

Carbohydrate sulfotransferase 2, 282-, Galactose/N-acetylglucosamine/N-acetylglucosamine 6-O-sulfotransferase 2, GST-2, N-acetylglucosamine 6-O-sulfotransferase 1, GlcNAc6ST-1, Gn6ST-1, CHST2, GN6ST

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**CHST2 Antibody (C-term) Blocking Peptide -  
Protein Information****Name** CHST2**Synonyms** GN6ST**Function**

Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the transfer of sulfate to position 6 of non-reducing N-acetylglucosamine (GlcNAc) residues

**CHST2 Antibody (C-term) Blocking Peptide  
- Background**

N-acetylglucosamine-6-O-sulfotransferases, such as CHST2, catalyze the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of a nonreducing N-acetylglucosamine (GlcNAc) residue (Uchimura et al., 1998 [PubMed9722682]).

**CHST2 Antibody (C-term) Blocking Peptide  
- References**

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010) Ross, C.J., et al. Nat. Genet. 41(12):1345-1349(2009) Desko, M.M., et al. Glycobiology 19(10):1068-1077(2009) Saito, A., et al. J. Hum. Genet. 54(6):317-323(2009) Kanoh, A., et al. Glycoconj. J. 23 (5-6), 453-460 (2006) :

within keratan-like structures on N-linked glycans and within mucin-associated glycans that can ultimately serve as SELL ligands. SELL ligands are present in high endothelial cells (HEVs) and play a central role in lymphocyte homing at sites of inflammation. Participates in biosynthesis of the SELL ligand sialyl 6-sulfo Lewis X and in lymphocyte homing to Peyer patches. Has no activity toward O-linked sugars. Its substrate specificity may be influenced by its subcellular location. Sulfates GlcNAc residues at terminal, non-reducing ends of oligosaccharide chains.

**Cellular Location**

Golgi apparatus, trans-Golgi network membrane; Single-pass type II membrane protein

**Tissue Location**

Widely expressed. Highly expressed in bone marrow, peripheral blood leukocytes, spleen, brain, spinal cord, ovary and placenta. Expressed by high endothelial cells (HEVs) and leukocytes

**CHST2 Antibody (C-term) Blocking Peptide  
- Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)