



NSE Blocking Peptide (C-term)

Synthetic peptide Catalog # BP19830b

Specification

NSE Blocking Peptide (C-term) - Product Information

Primary Accession Other Accession P09104

P07323, P17183,

NP 001966

NSE Blocking Peptide (C-term) - Additional Information

Gene ID 2026

Other Names

Gamma-enolase, 2-phospho-D-glycerate hydro-lyase, Enolase 2, Neural enolase, Neuron-specific enolase, NSE, ENO2

Target/Specificity

The synthetic peptide sequence is selected from aa 228-239 of HUMAN ENO2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

NSE Blocking Peptide (C-term) - Protein Information

Name ENO2

Function

Has neurotrophic and neuroprotective properties on a broad spectrum of central nervous system (CNS) neurons. Binds, in a

NSE Blocking Peptide (C-term) - Background

This gene encodes one of the three enclase isoenzymes found in mammals. This isoenzyme, a homodimer, is found in mature neurons and cells of neuronal origin. A switch from alpha enclase to gamma enclase occurs in neural tissue during development in rats and primates.

NSE Blocking Peptide (C-term) - References

Martins-de-Souza, D., et al. J Psychiatr Res 44(14):989-991(2010)
Mukhtarova, S.N. Georgian Med News 181, 49-54 (2010):
Planche, V., et al. Ann. Biol. Clin. (Paris) 68(2):239-242(2010)

Chaves, M.L., et al. J Neuroinflammation 7, 6 (2010):

Wijeyaratne, S.M., et al. Eur J Vasc Endovasc Surg 38(3):262-266(2009)





Tel: 858.875.1900 Fax: 858.622.0609

calcium- dependent manner, to cultured neocortical neurons and promotes cell survival (By similarity).

Cellular Location

Cytoplasm. Cell membrane. Note=Can translocate to the plasma membrane in either the homodimeric (alpha/alpha) or heterodimeric (alpha/gamma) form

Tissue Location

The alpha/alpha homodimer is expressed in embryo and in most adult tissues. The alpha/beta heterodimer and the beta/beta homodimer are found in striated muscle. and the alpha/gamma heterodimer and the gamma/gamma homodimer in neurons

NSE Blocking Peptide (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides