

PPP2CA Blocking Peptide (Center)

Synthetic peptide

Catalog # BP20181c

Specification**PPP2CA Blocking Peptide (Center) - Product Information**

Primary Accession [P67775](#)
Other Accession [P23696](#), [P63331](#),
[P67777](#), [P67776](#),
[P63330](#), [P67774](#),
[NP_002706.1](#)

PPP2CA Blocking Peptide (Center) - Additional Information**Gene ID** 5515**Other Names**

Serine/threonine-protein phosphatase 2A
catalytic subunit alpha isoform, PP2A-alpha,
Replication protein C, RP-C, PPP2CA

Target/Specificity

The synthetic peptide sequence is selected
from aa 103-116 of HUMAN PPP2CA

Format

Peptides are lyophilized in a solid powder
format. Peptides can be reconstituted in
solution using the appropriate buffer as
needed.

Storage

Maintain refrigerated at 2-8°C for up to 6
months. For long term storage store at
-20°C.

Precautions

This product is for research use only. Not
for use in diagnostic or therapeutic
procedures.

PPP2CA Blocking Peptide (Center) - Protein Information**Name** PPP2CA**Function**

PP2A is the major phosphatase for

PPP2CA Blocking Peptide (Center) - Background

This gene encodes the phosphatase 2A
catalytic subunit.
Protein phosphatase 2A is one of the four
major Ser/Thr
phosphatases, and it is implicated in the
negative control of cell
growth and division. It consists of a common
heteromeric core
enzyme, which is composed of a catalytic
subunit and a constant
regulatory subunit, that associates with a
variety of regulatory
subunits. This gene encodes an alpha isoform
of the catalytic
subunit.

PPP2CA Blocking Peptide (Center) - References

Shimada, M., et al. Hum. Genet.
128(4):433-441(2010)
Jayadeva, G., et al. J. Biol. Chem.
285(39):29863-29873(2010)
Pradhan, S., et al. J. Biol. Chem.
285(38):29059-29068(2010)
Schmitz, M.H., et al. Nat. Cell Biol.
12(9):886-893(2010)
Antony, R., et al. J. Biol. Chem.
285(24):18301-18308(2010)

microtubule-associated proteins (MAPs). PP2A can modulate the activity of phosphorylase B kinase casein kinase 2, mitogen-stimulated S6 kinase, and MAP-2 kinase. Cooperates with SGO2 to protect centromeric cohesin from separase-mediated cleavage in oocytes specifically during meiosis I (By similarity). Can dephosphorylate SV40 large T antigen and p53/TP53. Activates RAF1 by dephosphorylating it at 'Ser-259' (PubMed:10801873). Mediates dephosphorylation of WEE1, preventing its ubiquitin-mediated proteolysis, increasing WEE1 protein levels, and promoting the G2/M checkpoint (PubMed:33108758).

Cellular Location

Cytoplasm. Nucleus. Chromosome, centromere. Cytoplasm, cytoskeleton, spindle pole. Note=In prometaphase cells, but not in anaphase cells, localizes at centromeres. During mitosis, also found at spindle poles. Centromeric localization requires the presence of SGO2 (By similarity).

PPP2CA Blocking Peptide (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)